

11/13/00

11-14-00

A/Box/Seq

UTILITY PATENT APPLICATION TRANSMITTAL

(Only for new nonprovisional applications under 37 CFR 1.53(b))

Attorney Docket No. 35718/205458

First Inventor or Application Identifier: Johal et al.

Title of Invention: SORGHUM DWARFING GENES AND
METHODS OF USE

Express Mail Label No. EL613495598US

ADDRESS TO: ASSISTANT COMMISSIONER FOR PATENTS
BOX PATENT APPLICATION
WASHINGTON, DC 20231

Transmitted herewith for filing in the United States Patent Office is a patent application for:

Inventors: Gurmukh S. Johal and Dilbag S. Multani

1. ☒ The Filing Fee has been calculated as shown below:
2. ☐ Applicant claims Small Entity Status. See 37 CFR 1.27.

	No. Filed	No. Extra	Small Entity Rate	Fee 0	Large Entity Rate	Fee 0
BASIC FEE				\$0		\$710
TOTAL CLAIMS:	32 - 20 =	12	X 9 =	\$0	x 18 =	\$216
INDEP CLAIMS:	7 - 3 =	4	X 40 =	\$0	x 80 =	\$320
<input type="checkbox"/> MULTIPLE DEPENDENT CLAIMS PRESENTED			+135 =	\$	+270 =	\$
*If the difference in Column 1 is less than zero, enter "0" in Column 2.			TOTAL \$		TOTAL \$	1,246.00

The Commissioner is hereby authorized to credit overpayments or charge the following fees to Deposit Acct. No. 16-0605.

- a. ☒ Fees required under 37 CFR 1.16 (National filing fees).
- b. ☒ Fees required under 37 CFR 1.17 (National application processing fees).
- ☒ A check in the amount of \$ 1,246.00 for the filing fee is enclosed.
- ☐ The above filing fee will be paid along with Applicant(s) Response to the Notice to File Missing Parts.
3. ☒ Specification; Total Pages 59
4. ☐ _____ Sheets of Formal Drawing(s) (35 USC 113)
5. ☐ Declaration and Power of Attorney; [Total Pages _____]
 - a. ☐ Newly executed (original or copy)
 - b. ☐ Copy from a prior application (37 CFR 1.63(d))
(for continuation/divisional with Box 17 completed)
 - i. ☐ DELETION OF INVENTOR(S) Signed statement attached deleting inventor(s) named in the prior application, see 37 CFR 1.63(d)(2) & 1.33(b).
6. ☐ Application Data Sheet. See 37 CFR 1.76
7. ☐ CD-ROM or CD-R in duplicate, large table or Computer Program (Appendix)

8. Nucleotide and/or Amino Acid Sequence Submission (if applicable, all necessary)
- a. ☒ Computer Readable Copy (CRF)
 - b. ☒ Specification Sequence Listing on:
 - i. ☐ CD-ROM or CD-R (2 copies); or
 - ii. ☒ Paper
 - c. ☒ Statement verifying identity of above copies

ACCOMPANYING APPLICATION PARTS

9. ☐ Assignment Papers (cover sheet & document(s) (including a check for the \$40.00 fee)
10. ☐ 37 CFR 3.73(b) Statement (*when there is an assignee*); ☐ Power of Attorney
11. ☐ English Translation Document (*if applicable*)
12. ☐ Information Disclosure Statement (IDS)/PTO-1449; ___ Copies of IDS Citations
13. ☐ Preliminary Amendment
14. ☒ Return Receipt Postcard (MPEP 503) (*Should be specifically itemized*)
15. ☐ Certified Copy of Priority Document(s) (*if foreign priority is claimed*)
Foreign Priority is
16. ☐ Other: ___
17. **If a CONTINUING APPLICATION**, check appropriate box and supply the requisite information below and in a preliminary amendment, or in an Application Data Sheet under 37 CF 1.76:
☐ Continuation ☐ Divisional ☐ Continuation in Part (CIP)
of prior Application No: ___; Filed ___

Prior Application Information: Examiner

Group/Art Unit:

For CONTINUATION or DIVISIONAL APPS only: The entire disclosure of the prior application, from which an oath or declaration is supplied under Box 5b, is considered a part of the disclosure of the accompanying continuation or divisional application and is hereby incorporated by reference. The incorporation can only be relied upon when a portion has been inadvertently omitted from the submitted application parts.

18. **CORRESPONDENCE ADDRESS**

Customer Number or Bar Code Label **000826**

Attention Of: W. Murray Spruill

Signature: _____

Attorney/Agent of Record: W. Murray Spruill

Attorney/Agent Registration No. 32,943

Tel Raleigh Office (919) 420-2200

Fax Raleigh Office (919) 420-2260

ALSTON & BIRD LLP

P.O. Drawer 34009

Charlotte NC 28234-4009

"Express Mail" mailing label number EL613495598US

Date of Deposit November 13, 2000

I hereby certify that this paper or fee is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to Box Patent Application, Assistant Commissioner For Patents, Washington, DC 20231.

Sheree T. Edington

RTA01/2086289v1

SORGHUM DWARFING GENES AND METHODS OF USE

CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims the benefit of U.S. Provisional Application No. 60/165,176, filed November 12, 1999.

FIELD OF THE INVENTION

The present invention relates to the genetic manipulation of organisms, particularly plants, with genes that control growth and development. The invention further relates to genes that control growth, including homologues and mutant forms, the proteins encoded therefrom and plants transformed with these genes.

BACKGROUND OF THE INVENTION

Dwarf plants have had a major impact on agriculture. Dwarf varieties of wheat are widely used in North America due to both reduced potential for lodging and high yields. Dwarf fruit trees are also extensively used and allow farmers to produce more fruit per acre thereby increasing economic yield potential. There are other benefits that may be realized from the use of dwarf crop plants and dwarf fruit trees including reductions in the amounts of pesticides and fertilizers required, higher planting densities and reduced labor costs.

In view of the current trends of both increasing human population and the decreasing land area suitable for agriculture, increasing agricultural productivity is, and will continue to be, a challenge of paramount importance. Dwarf crop plants and fruit trees have been and will continue to be important components of our agricultural

production system. Increased usage of dwarf crop plants and dwarf fruit trees may help to meet the agricultural production demands of the future. However, commercially acceptable dwarf varieties are not available for all crops.

In addition to the use of dwarf plants to control plant height, synthetic chemicals are routinely applied to certain economically important plant species to reduce growth. Plant growth regulators known as growth retardants are used to reduce stem elongation in a variety of crops including cotton, grape vines, fruit trees, peanuts, wheat and ornamentals such as azaleas, chrysanthemums, hydrangeas, poinsettias and many bedding plants. All of the commonly used growth retardants are inhibitors of gibberellin biosynthesis and limit stem or shoot growth by reducing elongation. In the United States, the most widely used growth retardant is mepiquat chloride, which is registered for use on cotton. Benefits attributed to the use of mepiquat chloride on cotton include increased yield, improved defoliation, improved stress tolerance, more uniform crop maturity and the ability to harvest earlier. Previously, the growth retardant daminozide was registered for use in the United States on apples, grapes and peanuts under the trademarks ALAR and KYLAR but was removed from use on food crops due to human health concerns. Despite the demands of agricultural producers for a product to replace daminozide, there are no growth retardants registered for use on grapes, fruit trees and peanuts in the United States. Daminozide, however, is still widely used on certain non-food, plant species.

Uncovering the molecular mechanisms that control plant growth processes such as cell division and cell elongation will likely aid in the development of new plant varieties with reduced stature and new methods for reducing plant growth. Such new plant varieties and methods may provide both farmers and horticulturists with environmentally benign alternatives to the use of synthetic growth-retarding chemicals.

Elongation of plant cells and organs is one of the most critical parameters of plant growth and development. Regulation of this trait in plants, however, is a fairly complicated process, as both external and internal factors influence it. The most important external stimulus is light, with its normally repressible or negative effect on cell elongation (Quail, P.H. (1995) *Science* 268:675-680; Kende *et al.* (1997) *Plant Cell* 9:1197-1210). The internal control of cell elongation is mediated by a number of chemicals, normally referred to as plant growth regulators or hormones (Kende *et al.*

(1997) *Plant Cell* 9:1197-1210). Among the classical plant hormones, auxins and gibberellins (GAs) both promote cell elongation whereas cytokinins and abscisic acid each have been shown to have a negative effect on cell elongation (Kende *et al.* (1997) *Plant Cell* 9:1197-1210). Recently, another class of plant growth regulators, named
5 brassinosteroids, has been identified that also dramatically promote plant growth (Yokota, T. (1997) *Trends Plant Sci.* 2:137-143; Azpiroz *et al.* (1998) *Plant Cell* 10:219-230; Choe *et al.* (1998) *Plant Cell* 10:231-243). However, the mechanisms by which plant hormones act, either singly or in concert, to control cell elongation remains unclear.

One way to gain an understanding of mechanisms that mediate cell elongation is
10 to study mutants in which this aspect of plant growth is compromised (Klee *et al.* (1991) *Annu. Rev. Plant Physiol. Plant Mol. Biol.* 42:529-551). Numerous such mutants have been identified across most plant species, including maize, in which more than 25 single-gene mutations that affect plant stature have been characterized (Coe *et al.* (1988) In: *Corn & Corn Improvement*, G. F. Sprague (Ed.) Madison, WI ; Sheridan, W.F. (1988)
15 *Annu. Rev. Genet.* 22:353-385). These dwarf mutants are considered to be GA related, mainly because GA is the only phytohormone whose role in regulating height in maize has been convincingly established (Phinney *et al.* (1985) *Curr. Top. Plant Biochem. Physiol.* 4:67-74; Fujioka *et al.* (1988) *Proc. Natl. Acad. Sci. USA* 85:9031-9035). Both types of mutants, GA responsive and GA non-responsive, have been found in this
20 collection of maize mutants. While genes for a number of GA-responsive mutants have been cloned and found to be involved in GA biosynthesis (Bensen *et al.* (1995) *Plant Cell* 7:75-84; Winkler *et al.* (1995) *Plant Cell* 7:1307-1317), nothing is known about the nature of defects in GA non-responsive maize mutants.

One type of GA non-responsive dwarf mutants that have received much attention
25 from maize geneticists and breeders is called brachytic. These dwarfs are characterized by internodes of substantially reduced length, relative to wild-type, without having any effect on the size or number of other organs, including the leaves, ear and tassel (Kempton, J.H. (1920) *J. Hered.* 11:111-115). There are three known brachytic mutations in maize, *br1*, *br2* and *br3*, all of which are recessive (Coe *et al.* (1988) In:
30 *Corn & Corn Improvement*, G. F. Sprague (Ed.) Madison, WI; Sheridan, W.F. (1988) *Annu. Rev. Genet.* 22:353-385). Because of the commercial interest in *br2* for enhancing

plant productivity (Pendleton *et al.* (1961) *Crop Sci.* 1:433-435; Duvick, D.N. (1977) *Maydica* 22:187-196; Djisbar *et al.* (1987) *Maydica* 32:107-123; Russel, W.A. (1991) *Adv. Agron.* 46:245-298), this dwarf has been characterized the most. Depending on the genetic background, plants homozygous recessive for *br2* are 30-70% shorter than their normal sibs. This reduction in plant height is exclusively due to a reduction of the length of stalk (stem) internodes. In addition to being dwarf, *br2* mutants grown under greenhouse conditions often suffer from buggy whip, a disease-like condition in which the unfurling leaves in the whorl undergo necrosis and stay stuck together. This condition often results in the death of the growing tip of the plant.

Although the dwarfing trait in maize has been extensively studied both genetically and molecularly, it has yet to be exploited successfully in breeding efforts in this crop plant. In contrast, dwarf mutants of sorghum have contributed significantly to the development of modern day cultivars. Sorghum and maize are both members of the grass (Poaceae or Gramineae) family and thus share many characteristics including genomic organization and plant body form. Out of the four dwarfing mutations exploited in sorghum, *dw3*, whose dwarfing phenotype looks very similar to that of *br2* in maize, appears to be the most prominent. However, the only *dw3* allele (*dw3-ref*) available thus far has a serious problem which limits its agronomic value. The dwarf phenotype associated with the *dw3* allele is unstable, with a reversion frequency to wild-type (tall) as high as about 1% in certain genetic backgrounds. The instability of this dwarf phenotype, the mechanism of which has eluded sorghum geneticists thus far, not only continues to embarrass sorghum breeders, but also sometimes leads to the rejection of an otherwise promising inbred or hybrid.

To keep up with the demand for increased agricultural production, new targets are needed for genetically engineering agricultural plants for the improvement of agronomic characteristics. Elucidating the molecular mechanisms of cell division and elongation will provide new targets for agricultural scientists to manipulate.

SUMMARY OF THE INVENTION

Compositions and methods for expressing genes encoding P-glycoproteins in plants are provided. The compositions comprise nucleotide sequences encoding P-

glycoproteins, particularly P-glycoproteins that control plant growth. The compositions further comprise nucleotide sequences of the *Dw3* gene of sorghum. The sequences of the invention are useful in transforming plants for tissue-preferred or constitutive expression of P-glycoproteins and for isolating homologous nucleotide molecules that encode P-glycoproteins.. Such sequences find use in methods for controlling the growth of organisms, particularly stem growth in plants. The sequences of the invention also find use in methods of enhancing the resistance of plants to pathogens.

The invention further encompasses methods for isolating nucleotide molecules that are capable of controlling the growth of plants. Such methods find use in the isolation of genes involved in plant growth processes.

Methods are provided for identifying plants that possess a mutant allele that is capable of conferring a stable mutant phenotype on an organism. Such methods find use in agriculture, particularly in the breeding of dwarf crop plants, particularly dwarf sorghum plants.

Expression cassettes comprising the sequences of the invention are provided. Additionally provided are transformed plants, plant tissues, plant cells and seeds thereof. Isolated proteins encoded by the nucleotide sequences of the invention are provided.

DETAILED DESCRIPTION OF THE INVENTION

The present invention is drawn to compositions and methods for manipulating the growth of organisms. The methods involve transforming organisms with nucleotide sequences encoding P-glycoproteins. In particular, the nucleotide sequences are useful for controlling stem growth in plants. Thus, transformed plants, plant cells, plant tissues and seeds are provided. Compositions are nucleic acids and proteins relating to P-glycoprotein or P-glycoprotein-like genes in plants. More particularly, nucleotide sequences of the *Dw3* gene of sorghum and the amino acid sequences of the proteins encoded thereby are disclosed. The sequences find use in the construction of expression vectors for subsequent transformation into plants of interest, as probes for the isolation of other P-glycoprotein-like genes, as molecular markers, and the like.

The present invention discloses the first unequivocal evidence of the involvement of multidrug-resistance-like P-glycoproteins in the control of growth and development in

an organism. Thus, it is recognized that any P-glycoprotein known in the art that affects growth and development can be used in the practice of the invention. For example, five other plant P-glycoproteins are known. See, for example Dudler *et al.* (1998) *Methods Enzym.* 292:162-173 (*Arabidopsis*), Davies *et al.* (1997) *Gene* 199:195-202 (Barley),
5 Wang *et al.* (1996) *Plant Mol. Biol.* 31:683-687 (Potato) and GenBank Acession Numbers Y10227 and Y15990 (both from *Arabidopsis*); herein incorporated by reference. These and other P-glycoprotein sequences may be tested for an effect on growth by methods such as, for example, transformation with antisense sequences and monitoring effects on progeny plants.

10 The present invention also discloses methods for identifying genes encoding multidrug-resistance-like P-glycoproteins that control the growth of an organism, particularly a plant. An example of the identification of such a gene is disclosed for the *Dw3* gene of sorghum. Also provided is a method for identifying an allele of a gene wherein the allele confers a stable dwarf phenotype on a plant. An embodiment of this
15 method involves identifying stable mutant alleles of the *Dw3* gene that confer a dwarf phenotype on sorghum plants.

Compositions of the invention include the native nucleotide sequences for P-glycoprotein genes, antisense sequences, as well as variants and fragments thereof. Particularly, the P-glycoprotein gene of the sorghum *Dw3* locus and the respective amino
20 acid sequence for the P-glycoproteins encoded thereby, as well as fragments and variants thereof are provided. The *Dw3* nucleotide sequences are set forth in SEQ ID NOS: 1-3 and 7-8. The nucleotide sequences or corresponding antisense sequences find use in modulating the expression of a P-glycoprotein in a plant or plant cell. That is, the coding sequences can be used to increase the expression while antisense sequences can be used
25 to decrease expression.

The sequences of the invention find use in methods of modifying the growth of an organism. In an embodiment of the invention, nucleotide sequences of the invention find use in methods of modifying plant growth. Toward this end, the sequences of the invention may be utilized in expression cassettes or nucleotide constructs operably linked
30 to any one of a variety of plant promoters. Aspects of plant growth that may be impacted by the methods of the invention include, but are not limited to, plant height; the size,

shape and number of cells and organs; cell division rate; cell elongation rate; the growth rate of the plant, its organs, tissues and cells; timing and location of organ initiation; life span; and the like.

The invention discloses methods for reducing plant growth which find use as alternatives to applying synthetic, growth-retarding chemicals to plants. These methods provide environmentally safe alternatives to traditional means of retarding stem elongation or growth with synthetic chemicals. Some embodiments of the invention make use of plants transformed with tissue-preferred promoters, particularly stem-preferred promoters, operably linked to nucleotide sequences encoding P-glycoproteins.

Methods are provided for reducing the growth of a plant. Such methods involve transforming plants with at least one nucleotide sequence of the invention. The nucleotide sequences may be used in either the sense or antisense orientation to suppress the level of an endogenous P-glycoprotein that controls the growth of a plant. By reducing the level in a plant of such a P-glycoprotein, particularly one that controls stem or stalk growth, a plant of reduced stature, a dwarf plant, may be achieved. Dwarf plants having improved agronomic characteristics can be obtained by these methods. Such improved agronomic characteristics include, but are not limited to, reduced potential for lodging, increased water-use efficiency, reduced life cycle, increased harvest efficiency and increased yield per unit area. The methods of the invention can eliminate the need to graft shoots of fruit trees on dwarfing rootstocks to produce dwarf fruit trees.

The methods of the invention find use in producing dwarf varieties of crop plants. In one embodiment of the invention, a dwarf Basmati rice plant is produced by transforming the plant with a nucleotide sequence encoding at least a portion of a P-glycoprotein that controls the growth of a plant. Basmati rice, known for its aromatic fragrance, slender, elongated grains, and relatively short cooking time, is the favorite type of rice of the majority of people in the Indian sub-continent. While commercially acceptable dwarf cultivars have been developed for other types of rice, previous attempts to produce commercially acceptable varieties of Basmati rice by traditional plant breeding methods have failed. While dwarf plants were obtained in such attempts, some of the distinctive grain characteristics that consumers expect in Basmati rice were not retained in the dwarf plants. The methods of the invention provide a means of making

dwarf Basmati rice plants that produce grain possessing the characteristics desired by consumers.

The desired dwarf Basmati rice plants are produced by transforming a non-dwarf Basmati rice plant with a nucleotide sequence of the invention operably linked to a promoter that drives expression in a plant. While the choice of promoter depends on the desired outcome, the preferred promoters are tissue-preferred promoters, particularly stem-preferred promoters. Through cosuppression (sense suppression) or antisense suppression, such plants produce reduced levels of at least one P-glycoprotein that controls the growth of the Basmati rice plant, particularly stem growth. Preferably, the nucleotide sequence encodes at least a portion of a P-glycoprotein that controls the growth of a plant. More preferably, the nucleotide sequence is selected from the group consisting of SEQ ID NOS: 1-3 and 7-8 or a nucleotide sequence that encodes the amino acid sequence set forth in SEQ ID NOS: 4 or 9. Most preferably, the nucleotide sequence is from a rice gene that is homologous to the sorghum gene, *Dw3*. Such a rice gene encodes a P-glycoprotein that controls the growth of the stem of the rice plant. The methods of the invention comprise transforming plants with the full-length nucleotide sequences of the invention, or any fragment or part thereof.

Methods for enhancing the resistance of plants to pathogens are provided. It is recognized that P-glycoproteins are involved in resistance mechanisms against pathogens. A mutant strain of the nematode, *Caenorhabditis elegans*, with deletions of two P-glycoprotein genes is substantially more susceptible to death than wild-type nematodes, when placed on a lawn of a *Pseudomonas aeruginosa* strain that is a pathogen of both plants and animals (Mahajan-Miklos *et al.* (1999) *Cell* 96:47-56). *Br2* is a maize gene that encodes a multidrug-resistance-like P-glycoprotein that controls plant growth, particularly stem growth (See U.S. Provisional Application Serial No. 60/164,886 entitled "Genes and Methods for Manipulation of Growth" filed November 12, 1999; herein incorporated by reference). Maize plants that are homozygous for the mutant allele, *br2*, display a dwarf stature, and under certain cultural conditions, can also display a phenotype known as "buggy whip" which mimics a bacterial pathogen-induced necrosis of the growing tip of a plant.

The methods for the enhancing resistance of plants to pathogens comprise transforming plants with the nucleotide sequences of the invention operably linked to promoters that drive expression in a plant. Such plants display enhanced resistance to pathogens, including bacteria, fungi, viruses, nematodes and insects. The methods find
5 use in agriculture for limiting the impact of plant pathogens on crop production and provide an alternative to the use of synthetic pesticides in controlling plant pathogens.

Also provided are methods for identifying a plant with a stable mutant phenotype. Such methods find use in agriculture, particularly in the development of improved crop plants. The methods relate to an insertion-induced, mutant phenotype. By “insertion-
10 induced, mutant phenotype” is intended a mutant phenotype that is due to the insertion of a nucleotide, or a sequence of nucleotides, into the sequence of a gene of interest. While the invention does not depend upon a particular genetic mechanism for such an insertion-induced mutant phenotype, the presence of such an insertion within a gene typically disrupts the normal wild-type function of the gene, or gene product thereof. While the
15 methods of the invention are not bound by any particular type of insertion, such an insertion may be due to, for example, the insertion of a transposon or transposable element, or the duplication of a nucleotide sequence such as those which are known to occur as a result of genetic recombination.

Preferably, such an insertion-induced phenotype is unstable from one generation
20 to the next. That is, self pollination of one or more like plants having the insertion-induced phenotype results in at least one individual from among the resulting progeny population that has reverted to the wild-type phenotype. More preferably, such phenotypic instability, from one generation to the next, is due to the loss of at least a portion of the insertion from the gene of interest and that such a loss results in at least one
25 progeny plant, which has reverted to a wild-type phenotype. The methods of the present invention involve identifying an individual with a stable mutant phenotype from among such progeny population.

To identify a plant possessing an allele of a gene that confers a stable mutant phenotype, genomic DNA from a mutant plant is analyzed to determine if at least one
30 copy of the gene of interest lacks the insertion, or at least a portion thereof. Generally, the mutant plant is selected from a population of progeny derived from the self

pollination of one or more plants having the insertion-induced, mutant phenotype. Typically, in a population of such progeny, wild-type revertants will also be observed, indicating that at least a portion of the insertion has excised from the gene of interest. The genomic DNA of the selected mutant plant can be isolated and analyzed for the
5 absence of all or a portion of the insertion by techniques known to those of ordinary skill in the art such as, for example, Southern blotting, restriction fragment length polymorphism (RFLP) analysis and DNA amplification by polymerase chain reaction (PCR). Once a mutant plant lacking a portion of the insertion is identified, the progeny of such a mutant plant can be monitored to verify phenotypic stability. If desired,
10 subsequent generations can also be monitored.

Also provided are plants having stable mutant phenotypes and nucleotide sequences of alleles of genes which are capable of conferring a stable mutant phenotype on a plant.

A method of the invention involves identifying a sorghum plant with a stable
15 dwarf phenotype. Such a sorghum plant possesses in its genome a stable mutant allele of the *Dw3* gene. Such a stable mutant allele is capable of conferring a stable dwarf phenotype on a sorghum plant and the nucleotide sequence of a fragment of such an allele is set forth in SEQ ID NO: 2. One method of the invention employs RFLP analysis utilizing Southern blotting with a probe derived from nucleotide sequences of maize *Br2*.
20 This method additionally involves PCR amplification and DNA sequence analysis to determine the nucleotide sequence of the stable mutant allele.

Methods are provided for identifying nucleotide sequences encoding gene products that control plant growth. Such gene products, like the DW3 protein, impact or modify the growth of a plant in detectable way by, for example, affecting characteristics
25 such as the height or shape of a cell, organ or the plant body itself, cell number, cell division rate or cell elongation rate, organ growth rate, appearance of reproductive structures, timing and location of organ initiation and the like. The methods of the invention are particularly directed toward nucleotide sequences which influence the height or stature of a plant. The nucleotide sequences of the invention find use in any
30 method known to those skilled in the art for identifying homologous sequences. Such

methods for identifying homologous sequences include PCR amplification, hybridization, Southern blotting, colony hybridization and the like.

An embodiment of the invention involves the use of PCR amplification to identify nucleotide sequences encoding gene products that control plant growth. Such PCR
5 amplification comprises the use of at least one oligonucleotide primer derived from a nucleotide sequence encoding of a gene encoding a multidrug-resistance-like P-glycoprotein. Preferably, such a nucleotide sequence is from a gene that encodes a P-glycoprotein that controls the growth of an organism, particularly a plant. More preferably, the nucleotide sequence is selected from the group consisting of SEQ ID
10 NOS: 1-3 and 7-8.

In another embodiment, oligonucleotide primers (SEQ ID NOS: 5-6) were prepared from the sequences of *Br2*. Such primers were used to PCR amplify *Dw3* from genomic DNA isolated from sorghum plants. Following DNA sequencing the identity of *Dw3* was revealed. In a similar manner, other homologues of both *Br2* and *Dw3* can be
15 identified using the same primers or other primers derived from any gene encoding a P-glycoprotein that controls the growth of an organism.

In still another exemplary embodiment of the invention, one or more nucleotide sequences set forth in SEQ ID NOS: 1-3 and 5-8 or a nucleotide sequence encoding the amino acid sequence set forth in SEQ ID. NO. 4 or 9 are used to design hybridization
20 probes or PCR primers to identify a gene in the genome of a Basmati rice plant that is homologous to the sorghum gene, *Dw3*. Preferably, such a gene, from a Basmati rice plant, encodes a P-glycoprotein. More preferably, such a gene encodes a P-glycoprotein that controls the growth of the Basmati rice plant. Most preferably, such a gene encodes a P-glycoprotein that controls the stem growth of the Basmati rice plant.

25 The P-glycoproteins of the invention encompass all polypeptides and nucleotide sequences encoding them that share substantial sequence identity to the sequences of the invention whether or not such polypeptides possess covalently attached carbohydrates or carbohydrate-containing chains.

By "control growth of an organism" is intended to include impacting, modifying,
30 modulating, affecting, increasing, and decreasing growth and growth-related processes of an organism. Such processes may influence any of a multitude of characteristics of an

organism including, but not limited to, cell size and shape, organism size and shape, cell division rate, cell enlargement rate, organ growth rate, onset of reproductive maturity and life span.

By “mutant phenotype” is intended any non-wild-type, non-typical or non-standard phenotype which occurs as a result of a genetic alteration in the genome of an organism. When used in reference to domesticated plants and animals, a “mutant phenotype” is any phenotype that is substantially different from the typical phenotype of the particular domesticated breed or cultivated variety from which the mutant phenotype arose.

By “mutant plant” is intended a plant having a mutant phenotype.

By “mutant allele” is intended an allele of a gene that is capable of causing a “mutant phenotype.”

By “dwarf” is intended atypically small. By “dwarf plant” is intended an atypically small plant. Generally, such a “dwarf plant” has a stature or height that is reduced from that of a typical plant by about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60% or greater. Generally, but not exclusively, such a dwarf plant is characterized by a reduced stem, stalk or trunk length when compared to the typical plant.

By “nucleotide molecule” is intended a molecule composed of nucleotides covalently bound to one another. Nucleotides include both ribonucleotides and deoxyribonucleotides. “Nucleotide molecule” encompasses single-stranded and double stranded forms of both DNA and RNA. “Nucleotide molecules” may be naturally occurring, synthetic or a combination of both. The linear arrangement of nucleotides in a “nucleotide molecule” is referred to as a “nucleotide sequence” and unless specified otherwise is presented herein from left to right corresponding to 5'-to-3' direction. Because of the complementary nature of the opposite strands of a double-stranded nucleotide molecule, a nucleotide sequence of the invention additionally encompasses its complementary antisense sequence.

Compositions of the invention include native nucleotide sequences for genes encoding multidrug-resistance-like-gene-encoded P-glycoproteins, homologues of multidrug-resistance-like-gene-encoded P-glycoproteins, antisense sequences, as well as fragments and variants and fragments thereof. In particular, the present invention

provides for isolated nucleic acid molecules comprising nucleotide sequences encoding the amino acid sequences shown in SEQ ID NOS: 4 and 9, or the nucleotide sequences encoding the DNA sequences deposited in a bacterial host as Patent Deposit No. PTA 2646. Further provided are polypeptides having an amino acid sequence encoded by a nucleic acid molecule described herein, for example those set forth in SEQ ID NOS: 3 and 8, respectively, those deposited in a bacterial host as Patent Deposit Nos. PTA 2646, and fragments and variants thereof.

Plasmids containing the nucleotide sequences of the invention were deposited with the Patent Depository of the American Type Culture Collection (ATCC), Manassas, Virginia, on November 1, 2000 and assigned Patent Deposit No PTA 2646. These deposits will be maintained under the terms of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purposes of Patent Procedure. These deposits were made merely as a convenience for those of skill in the art and are not an admission that a deposit is required under 35 U.S.C. §112.

The invention encompasses isolated or substantially purified nucleic acid or protein compositions. An "isolated" or "purified" nucleic acid molecule or protein, or biologically active portion thereof, is substantially free of other cellular material, or culture medium when produced by recombinant techniques, or substantially free of chemical precursors or other chemicals when chemically synthesized. Preferably, an "isolated" nucleic acid is free of sequences (preferably protein encoding sequences) that naturally flank the nucleic acid (i.e., sequences located at the 5' and 3' ends of the nucleic acid) in the genomic DNA of the organism from which the nucleic acid is derived. For example, in various embodiments, the isolated nucleic acid molecule can contain less than about 5 kb, 4 kb, 3 kb, 2 kb, 1 kb, 0.5 kb, or 0.1 kb of nucleotide sequences that naturally flank the nucleic acid molecule in genomic DNA of the cell from which the nucleic acid is derived. A protein that is substantially free of cellular material includes preparations of protein having less than about 30%, 20%, 10%, 5%, (by dry weight) of contaminating protein. When the protein of the invention or biologically active portion thereof is recombinantly produced, preferably culture medium represents less than about 30%, 20%, 10%, or 5% (by dry weight) of chemical precursors or non-protein-of-interest chemicals.

Fragments and variants of the disclosed nucleotide sequences and proteins encoded thereby are also encompassed by the present invention. By "fragment" is intended a portion of the nucleotide sequence or a portion of the amino acid sequence and hence protein encoded thereby. Fragments of a nucleotide sequence may encode protein fragments that retain biological activity of the native P-glycoprotein and hence retain one or more functions of the native P-glycoprotein such as, for example, transmembrane transporter activity and ATP binding. Alternatively, fragments of a nucleotide sequence that are useful as hybridization probes may or may not encode protein fragments retaining biological activity. Thus, fragments of a nucleotide sequence may range from at least about 20 nucleotides, about 50 nucleotides, about 100 nucleotides, and up to the full-length nucleotide sequence of the invention.

A fragment of a P-glycoprotein gene nucleotide sequence that encodes a biologically active portion of a P-glycoprotein of the invention will encode at least 15, 20, 25, 30, 50, 75, 100, 150, 200, 250, 300, 350, 400, 450, 500, 550, 600, 650, 700, 800, 900, 1,000, 1,100, 1,200, 1,300, or 1,400 contiguous amino acids, or up to the total number of amino acids present in a full-length P-glycoprotein of the invention (for example, 415 and 1,421 amino acids for SEQ ID NOS: 4 and 9). Fragments of a P-glycoprotein gene nucleotide sequence that are useful as hybridization probes for PCR primers generally need not encode a biologically active portion of a P-glycoprotein.

Thus, a fragment of a P-glycoprotein gene nucleotide sequence may encode a biologically active portion of a P-glycoprotein, or it may be a fragment that can be used as a hybridization probe or PCR primer using methods disclosed below. A biologically active portion of a P-glycoprotein can be prepared by isolating a portion of one of the P-glycoprotein gene nucleotide sequences of the invention, expressing the encoded portion of the P-glycoprotein e.g., by recombinant expression *in vitro*), and assessing the activity of the portion of the P-glycoprotein. Nucleic acid molecules that are fragments of a P-glycoprotein gene nucleotide sequence comprise at least 16, 20, 50, 75, 100, 150, 200, 300, 500, 700, 1,000, 1,200, 1,500, 2,000, 2,500, 3,000, 3,500, 4,000, 4,500, 5,000, 5,500, 6,000 nucleotides, or up to the number of nucleotides present in a full-length P-glycoprotein nucleotide sequence disclosed herein (for example, 2,139, 1,267, 1,261, 6,827, and 4213 nucleotides for SEQ ID NOS: 1-3, and 7-8, respectively).

By “variants” is intended substantially similar sequences. For nucleotide sequences, conservative variants include those sequences that, because of the degeneracy of the genetic code, encode the amino acid sequence of one of the P-glycoprotein polypeptides of the invention. Naturally occurring allelic variants such as these can be identified with the use of well-known molecular biology techniques, as, for example, with polymerase chain reaction (PCR) and hybridization techniques as outlined below. Variant nucleotide sequences also include synthetically derived nucleotide sequences, such as those generated, for example, by using site-directed mutagenesis but which still encode a P-glycoprotein protein of the invention. Generally, variants of a particular nucleotide sequence of the invention will have at least about 40%, 50%, 60%, 65%, 70%, generally at least about 75%, 80%, 85%, preferably at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, and more preferably at least about 98%, 99% or more sequence identity to that particular nucleotide sequence as determined by sequence alignment programs described elsewhere herein using default parameters.

By “variant” protein is intended a protein derived from the native protein by deletion (so-called truncation) or addition of one or more amino acids to the N-terminal and/or C-terminal end of the native protein; deletion or addition of one or more amino acids at one or more sites in the native protein; or substitution of one or more amino acids at one or more sites in the native protein. Variant proteins encompassed by the present invention are biologically active, that is they continue to possess the desired biological activity of the native protein, that is, transporter activity or ATP binding activity as described herein. Such variants may result from, for example, genetic polymorphism or from human manipulation. Biologically active variants of a native P-glycoprotein of the invention will have at least about 40%, 50%, 60%, 65%, 70%, generally at least about 75%, 80%, 85%, preferably at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, and more preferably at least about 98%, 99% or more sequence identity to the amino acid sequence for the native protein as determined by sequence alignment programs described elsewhere herein using default parameters. A biologically active variant of a protein of the invention may differ from that protein by as few as 1-15 amino acid residues, as few as 1-10, such as 6-10, as few as 5, as few as 4, 3, 2, or even 1 amino acid residue.

The proteins of the invention may be altered in various ways including amino acid substitutions, deletions, truncations, and insertions. Methods for such manipulations are generally known in the art. For example, amino acid sequence variants of the P-glycoproteins can be prepared by mutations in the DNA. Methods for mutagenesis and nucleotide sequence alterations are well known in the art. See, for example, Kunkel (1985) *Proc. Natl. Acad. Sci. USA* 82:488-492; Kunkel *et al.* (1987) *Methods in Enzymol.* 154:367-382; US Patent No. 4,873,192; Walker and Gaastra, eds. (1983) *Techniques in Molecular Biology* (MacMillan Publishing Company, New York) and the references cited therein. Guidance as to appropriate amino acid substitutions that do not affect biological activity of the protein of interest may be found in the model of Dayhoff *et al.* (1978) *Atlas of Protein Sequence and Structure* (Natl. Biomed. Res. Found., Washington, D.C.), herein incorporated by reference. Conservative substitutions, such as exchanging one amino acid with another having similar properties, may be preferred.

Thus, the genes and nucleotide sequences of the invention include both the naturally occurring sequences as well as mutant forms. Likewise, the proteins of the invention encompass both naturally occurring proteins as well as variations and modified forms thereof. Such variants will continue to possess the desired transporter activity. Obviously, the mutations that will be made in the DNA encoding the variant must not place the sequence out of reading frame and preferably will not create complementary regions that could produce secondary mRNA structure. See, EP Patent Application Publication No. 75,444.

The deletions, insertions, and substitutions of the protein sequences encompassed herein are not expected to produce radical changes in the characteristics of the protein. However, when it is difficult to predict the exact effect of the substitution, deletion, or insertion in advance of doing so, one skilled in the art will appreciate that the effect will be evaluated by routine screening assays.

Variant nucleotide sequences and proteins also encompass nucleotide sequences and proteins derived from a mutagenic and recombinogenic procedure such as DNA shuffling. With such a procedure, one or more different P-glycoprotein coding sequences can be manipulated to create a variant nucleotide sequence encoding a variant P-glycoprotein possessing the desired properties. In this manner, libraries of recombinant

polynucleotides are generated from a population of related sequence polynucleotides comprising sequence regions that have substantial sequence identity and can be homologously recombined *in vitro* or *in vivo*. For example, using this approach, sequence motifs encoding a domain of interest may be shuffled between the P-glycoprotein gene of the invention and other known P-glycoprotein genes to obtain a new gene coding for a protein with an improved property of interest, such as an increased K_m in the case of an enzyme. Strategies for such DNA shuffling are known in the art. See, for example, Stemmer (1994) *Proc. Natl. Acad. Sci. USA* 91:10747-10751; Stemmer (1994) *Nature* 370:389-391; Cramer *et al.* (1997) *Nature Biotech.* 15:436-438; Moore *et al.* (1997) *J. Mol. Biol.* 272:336-347; Zhang *et al.* (1997) *Proc. Natl. Acad. Sci. USA* 94:4504-4509; Cramer *et al.* (1998) *Nature* 391:288-291; and U.S. Patent Nos. 5,605,793 and 5,837,458.

The nucleotide sequences of the invention can be used to isolate corresponding sequences from other organisms, particularly other plants, more particularly other monocots. In this manner, methods such as PCR, hybridization, and the like can be used to identify such sequences based on their sequence homology to the sequences set forth herein. Sequences isolated based on their sequence identity to the entire sequences set forth herein or to fragments thereof are encompassed by the present invention. Such sequences include sequences that are orthologs of the disclosed sequences. By “orthologs” is intended genes derived from a common ancestral gene and which are found in different species as a result of speciation. Genes found in different species are considered orthologs when their nucleotide sequences and/or their encoded protein sequences share substantial identity as defined elsewhere herein. Functions of orthologs are often highly conserved among species.

In a PCR approach, oligonucleotide primers can be designed for use in PCR reactions to amplify corresponding DNA sequences from cDNA or genomic DNA extracted from any organism of interest. Methods for designing PCR primers and PCR cloning are generally known in the art and are disclosed in Sambrook *et al.* (1989) *Molecular Cloning: A Laboratory Manual* (2nd ed., Cold Spring Harbor Laboratory Press, Plainview, New York). See also Innis *et al.*, eds. (1990) *PCR Protocols: A Guide to Methods and Applications* (Academic Press, New York); Innis and Gelfand, eds.

(1995) *PCR Strategies* (Academic Press, New York); and Innis and Gelfand, eds. (1999) *PCR Methods Manual* (Academic Press, New York). Known methods of PCR include, but are not limited to, methods using paired primers, nested primers, single specific primers, degenerate primers, gene-specific primers, vector-specific primers, partially-mismatched primers, and the like.

In hybridization techniques, all or part of a known nucleotide sequence is used as a probe that selectively hybridizes to other corresponding nucleotide sequences present in a population of cloned genomic DNA fragments or cDNA fragments (*i.e.*, genomic or cDNA libraries) from a chosen organism. The hybridization probes may be genomic DNA fragments, cDNA fragments, RNA fragments, or other oligonucleotides, and may be labeled with a detectable group such as ³²P, or any other detectable marker. Thus, for example, probes for hybridization can be made by labeling synthetic oligonucleotides based on the P-glycoprotein gene nucleotide sequences of the invention. Methods for preparation of probes for hybridization and for construction of cDNA and genomic libraries are generally known in the art and are disclosed in Sambrook *et al.* (1989) *Molecular Cloning: A Laboratory Manual* (2nd ed., Cold Spring Harbor Laboratory Press, Plainview, New York).

For example, the entire *Dw3* sequence disclosed herein, or one or more portions thereof, may be used as a probe capable of specifically hybridizing to corresponding P-glycoprotein gene sequences and messenger RNAs. To achieve specific hybridization under a variety of conditions, such probes include sequences that are unique among P-glycoprotein gene sequences and are preferably at least about 10 nucleotides in length, and most preferably at least about 20 nucleotides in length. Such probes may be used to amplify corresponding P-glycoprotein gene sequences from a chosen plant by PCR. This technique may be used to isolate additional coding sequences from a desired plant or as a diagnostic assay to determine the presence of coding sequences in a plant. Hybridization techniques include hybridization screening of plated DNA libraries (either plaques or colonies; see, for example, Sambrook *et al.* (1989) *Molecular Cloning: A Laboratory Manual* (2nd ed., Cold Spring Harbor Laboratory Press, Plainview, New York).

Hybridization of such sequences may be carried out under stringent conditions. By “stringent conditions” or “stringent hybridization conditions” is intended conditions

under which a probe will hybridize to its target sequence to a detectably greater degree than to other sequences (e.g., at least two-fold over background). Stringent conditions are sequence-dependent and will be different in different circumstances. By controlling the stringency of the hybridization and/or washing conditions, target sequences that are 100% complementary to the probe can be identified (homologous probing).

Alternatively, stringency conditions can be adjusted to allow some mismatching in sequences so that lower degrees of similarity are detected (heterologous probing). Generally, a probe is less than about 1000 nucleotides in length, preferably less than 500 nucleotides in length.

Typically, stringent conditions will be those in which the salt concentration is less than about 1.5 M Na ion, typically about 0.01 to 1.0 M Na ion concentration (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30°C for short probes (e.g., 10 to 50 nucleotides) and at least about 60°C for long probes (e.g., greater than 50 nucleotides). Stringent conditions may also be achieved with the addition of destabilizing agents such as formamide. Exemplary low stringency conditions include hybridization with a buffer solution of 30 to 35% formamide, 1 M NaCl, 1% SDS (sodium dodecyl sulphate) at 37°C, and a wash in 1X to 2X SSC (20X SSC = 3.0 M NaCl/0.3 M trisodium citrate) at 50 to 55°C. Exemplary moderate stringency conditions include hybridization in 40 to 45% formamide, 1 M NaCl, 1% SDS at 37°C, and a wash in 0.5X to 1X SSC at 55 to 60°C. Exemplary high stringency conditions include hybridization in 50% formamide, 1 M NaCl, 1% SDS at 37°C, and a wash in 0.1X SSC at 60 to 65°C. The duration of hybridization is generally less than about 24 hours, usually about 4 to about 12 hours.

Specificity is typically the function of post-hybridization washes, the critical factors being the ionic strength and temperature of the final wash solution. For DNA-DNA hybrids, the T_m can be approximated from the equation of Meinkoth and Wahl (1984) *Anal. Biochem.* 138:267-284: $T_m = 81.5^\circ\text{C} + 16.6 (\log M) + 0.41 (\%GC) - 0.61 (\% \text{ form}) - 500/L$; where M is the molarity of monovalent cations, %GC is the percentage of guanosine and cytosine nucleotides in the DNA, % form is the percentage of formamide in the hybridization solution, and L is the length of the hybrid in base pairs. The T_m is the temperature (under defined ionic strength and pH) at which 50% of a

complementary target sequence hybridizes to a perfectly matched probe. T_m is reduced by about 1°C for each 1% of mismatching; thus, T_m , hybridization, and/or wash conditions can be adjusted to hybridize to sequences of the desired identity. For example, if sequences with >90% identity are sought, the T_m can be decreased 10°C. Generally, stringent conditions are selected to be about 5°C lower than the thermal melting point (T_m) for the specific sequence and its complement at a defined ionic strength and pH. However, severely stringent conditions can utilize a hybridization and/or wash at 1, 2, 3, or 4°C lower than the thermal melting point (T_m); moderately stringent conditions can utilize a hybridization and/or wash at 6, 7, 8, 9, or 10°C lower than the thermal melting point (T_m); low stringency conditions can utilize a hybridization and/or wash at 11, 12, 13, 14, 15, or 20°C lower than the thermal melting point (T_m). Using the equation, hybridization and wash compositions, and desired T_m , those of ordinary skill will understand that variations in the stringency of hybridization and/or wash solutions are inherently described. If the desired degree of mismatching results in a T_m of less than 45°C (aqueous solution) or 32°C (formamide solution), it is preferred to increase the SSC concentration so that a higher temperature can be used. An extensive guide to the hybridization of nucleic acids is found in Tijssen (1993) *Laboratory Techniques in Biochemistry and Molecular Biology—Hybridization with Nucleic Acid Probes*, Part I, Chapter 2 (Elsevier, New York); and Ausubel *et al.*, eds. (1995) *Current Protocols in Molecular Biology*, Chapter 2 (Greene Publishing and Wiley-Interscience, New York). See Sambrook *et al.* (1989) *Molecular Cloning: A Laboratory Manual* (2nd ed., Cold Spring Harbor Laboratory Press, Plainview, New York).

Thus, isolated sequences that encode for P-glycoproteins and which hybridize under stringent conditions to the to the P-glycoprotein gene sequences disclosed herein, or to fragments thereof, are encompassed by the present invention. Such sequences will be at least about 70% to 75%, about 80% to 85%, and even 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more homologous with the disclosed sequences. That is, the sequence identity of sequences may range, sharing at least about 70% to 75%, about 80% to 85%, and even at least about 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more sequence identity.

The following terms are used to describe the sequence relationships between two or more nucleic acids or polynucleotides: (a) "reference sequence", (b) "comparison window", (c) "sequence identity", (d) "percentage of sequence identity", and (e) "substantial identity."

5 (a) As used herein, "reference sequence" is a defined sequence used as a basis for sequence comparison. A reference sequence may be a subset or the entirety of a specified sequence; for example, as a segment of a full-length cDNA or gene sequence, or the complete cDNA or gene sequence.

10 (b) As used herein, "comparison window" makes reference to a contiguous and specified segment of a polynucleotide sequence, wherein the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. Generally, the comparison window is at least 20 contiguous nucleotides in length, and optionally can be 30, 40, 50, 100, or longer. Those
15 of skill in the art understand that to avoid a high similarity to a reference sequence due to inclusion of gaps in the polynucleotide sequence a gap penalty is typically introduced and is subtracted from the number of matches.

Methods of alignment of sequences for comparison are well known in the art. Thus, the determination of percent identity between any two sequences can be
20 accomplished using a mathematical algorithm. Non-limiting examples of such mathematical algorithms are the algorithm of Myers and Miller (1988) CABIOS 4:11-17; the local homology algorithm of Smith et al. (1981) Adv. Appl. Math. 2:482; the homology alignment algorithm of Needleman and Wunsch (1970) J. Mol. Biol. 48:443-453; the search-for-similarity-method of Pearson and Lipman (1988) Proc. Natl. Acad.
25 Sci. 85:2444-2448; the algorithm of Karlin and Altschul (1990) Proc. Natl. Acad. Sci. USA 87:2264, modified as in Karlin and Altschul (1993) Proc. Natl. Acad. Sci. USA 90:5873-5877.

Computer implementations of these mathematical algorithms can be utilized for comparison of sequences to determine sequence identity. Such implementations include,
30 but are not limited to: CLUSTAL in the PC/Gene program (available from Intelligenetics, Mountain View, California); the ALIGN program (Version 2.0) and GAP,

BESTFIT, BLAST, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Version 8 (available from Genetics Computer Group (GCG), 575 Science Drive, Madison, Wisconsin, USA). Alignments using these programs can be performed using the default parameters. The CLUSTAL program is well described by Higgins et al. (1988) *Gene* 73:237-244 (1988); Higgins et al. (1989) *CABIOS* 5:151-153; Corpet et al. (1988) *Nucleic Acids Res.* 16:10881-90; Huang et al. (1992) *CABIOS* 8:155-65; and Pearson et al. (1994) *Meth. Mol. Biol.* 24:307-331. The ALIGN program is based on the algorithm of Myers and Miller (1988) *supra*. A PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4 can be used with the ALIGN program when comparing amino acid sequences. The BLAST programs of Altschul et al (1990) *J. Mol. Biol.* 215:403 are based on the algorithm of Karlin and Altschul (1990) *supra*. BLAST nucleotide searches can be performed with the BLASTN program, score = 100, wordlength = 12, to obtain nucleotide sequences homologous to a nucleotide sequence encoding a protein of the invention. BLAST protein searches can be performed with the BLASTX program, score = 50, wordlength = 3, to obtain amino acid sequences homologous to a protein or polypeptide of the invention. To obtain gapped alignments for comparison purposes, Gapped BLAST (in BLAST 2.0) can be utilized as described in Altschul et al. (1997) *Nucleic Acids Res.* 25:3389. Alternatively, PSI-BLAST (in BLAST 2.0) can be used to perform an iterated search that detects distant relationships between molecules. See Altschul et al. (1997) *supra*. When utilizing BLAST, Gapped BLAST, PSI-BLAST, the default parameters of the respective programs (e.g., BLASTN for nucleotide sequences, BLASTX for proteins) can be used. See <http://www.ncbi.nlm.nih.gov>. Alignment may also be performed manually by inspection.

Unless otherwise stated, sequence identity/similarity values provided herein refer to the value obtained using GAP Version 10 using the following parameters: % identity using GAP Weight of 50 and Length Weight of 3; % similarity using Gap Weight of 12 and Length Weight of 4, or any equivalent program. By “equivalent program” is intended any sequence comparison program that, for any two sequences in question, generates an alignment having identical nucleotide or amino acid residue matches and an identical percent sequence identity when compared to the corresponding alignment generated by the preferred program.

GAP uses the algorithm of Needleman and Wunsch (1970) J. Mol. Biol. 48: 443-453, to find the alignment of two complete sequences that maximizes the number of matches and minimizes the number of gaps. GAP considers all possible alignments and gap positions and creates the alignment with the largest
5 number of matched bases and the fewest gaps. It allows for the provision of a gap creation penalty and a gap extension penalty in units of matched bases. GAP must make a profit of gap creation penalty number of matches for each gap it inserts. If a gap extension penalty greater than zero is chosen, GAP must, in addition, make a profit for each gap inserted of the length of the gap times the gap extension penalty. Default gap
10 creation penalty values and gap extension penalty values in Version 10 of the Wisconsin Genetics Software Package for protein sequences are 8 and 2, respectively. For nucleotide sequences the default gap creation penalty is 50 while the default gap extension penalty is 3. The gap creation and gap extension penalties can be expressed as an integer selected from the group of integers consisting of from 0 to 200. Thus, for
15 example, the gap creation and gap extension penalties can be 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65 or greater.

GAP presents one member of the family of best alignments. There may be many members of this family, but no other member has a better quality. GAP displays four figures of merit for alignments: Quality, Ratio, Identity, and Similarity. The Quality is
20 the metric maximized in order to align the sequences. Ratio is the quality divided by the number of bases in the shorter segment. Percent Identity is the percent of the symbols that actually match. Percent Similarity is the percent of the symbols that are similar. Symbols that are across from gaps are ignored. A similarity is scored when the scoring matrix value for a pair of symbols is greater than or equal to 0.50, the similarity
25 threshold. The scoring matrix used in Version 10 of the Wisconsin Genetics Software Package is BLOSUM62 (see Henikoff and Henikoff (1989) Proc. Natl. Acad. Sci. USA 89:10915).

(c) As used herein, "sequence identity" or "identity" in the context of two nucleic acid or polypeptide sequences makes reference to the residues in the two
30 sequences that are the same when aligned for maximum correspondence over a specified comparison window. When percentage of sequence identity is used in reference to

proteins it is recognized that residue positions which are not identical often differ by conservative amino acid substitutions, where amino acid residues are substituted for other amino acid residues with similar chemical properties (e.g., charge or hydrophobicity) and therefore do not change the functional properties of the molecule. When sequences differ in conservative substitutions, the percent sequence identity may be adjusted upwards to correct for the conservative nature of the substitution. Sequences that differ by such conservative substitutions are said to have "sequence similarity" or "similarity." Means for making this adjustment are well known to those of skill in the art. Typically this involves scoring a conservative substitution as a partial rather than a full mismatch, thereby increasing the percentage sequence identity. Thus, for example, where an identical amino acid is given a score of 1 and a non-conservative substitution is given a score of zero, a conservative substitution is given a score between zero and 1. The scoring of conservative substitutions is calculated, e.g., as implemented in the program PC/GENE (Intelligenetics, Mountain View, California).

(d) As used herein, "percentage of sequence identity" means the value determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison, and multiplying the result by 100 to yield the percentage of sequence identity.

(e)(i) The term "substantial identity" of polynucleotide sequences means that a polynucleotide comprises a sequence that has at least 70% sequence identity, preferably at least 80%, more preferably at least 90%, and most preferably at least 95%, compared to a reference sequence using one of the alignment programs described using standard parameters. One of skill in the art will recognize that these values can be appropriately adjusted to determine corresponding identity of proteins encoded by two nucleotide sequences by taking into account codon degeneracy, amino acid similarity, reading frame

positioning, and the like. Substantial identity of amino acid sequences for these purposes normally means sequence identity of at least 60%, more preferably at least 70%, 80%, 90%, and most preferably at least 95%.

Another indication that nucleotide sequences are substantially identical is if two molecules hybridize to each other under stringent conditions. Generally, stringent conditions are selected to be about 5°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. However, stringent conditions encompass temperatures in the range of about 1°C to about 20°C lower than the T_m , depending upon the desired degree of stringency as otherwise qualified herein. Nucleic acids that do not hybridize to each other under stringent conditions are still substantially identical if the polypeptides they encode are substantially identical. This may occur, e.g., when a copy of a nucleic acid is created using the maximum codon degeneracy permitted by the genetic code. One indication that two nucleic acid sequences are substantially identical is when the polypeptide encoded by the first nucleic acid is immunologically cross reactive with the polypeptide encoded by the second nucleic acid.

(e)(ii) The term “substantial identity” in the context of a peptide indicates that a peptide comprises a sequence with at least 70% sequence identity to a reference sequence, preferably 80%, more preferably 85%, most preferably at least 90% or 95% sequence identity to the reference sequence over a specified comparison window. Preferably, optimal alignment is conducted using the homology alignment algorithm of Needleman *et al.* (1970) *J. Mol. Biol.* 48:443. An indication that two peptide sequences are substantially identical is that one peptide is immunologically reactive with antibodies raised against the second peptide. Thus, a peptide is substantially identical to a second peptide, for example, where the two peptides differ only by a conservative substitution. Peptides that are “substantially similar” share sequences as noted above except that residue positions that are not identical may differ by conservative amino acid changes.

The use of the term “nucleotide constructs” herein is not intended to limit the present invention to nucleotide constructs comprising DNA. Those of ordinary skill in the art will recognize that nucleotide constructs, particularly polynucleotides and oligonucleotides, comprised of ribonucleotides and combinations of ribonucleotides and deoxyribonucleotides may also be employed in the methods disclosed herein. Thus, the

nucleotide constructs of the present invention encompass all nucleotide constructs that can be employed in the methods of the present invention for transforming plants including, but not limited to, those comprised of deoxyribonucleotides, ribonucleotides, and combinations thereof. Such deoxyribonucleotides and ribonucleotides include both naturally occurring molecules and synthetic analogues. The nucleotide constructs of the invention also encompass all forms of nucleotide constructs including, but not limited to, single-stranded forms, double-stranded forms, hairpins, stem-and-loop structures, and the like.

Furthermore, it is recognized that the methods of the invention may employ a nucleotide construct that is capable of directing, in a transformed plant, the expression of at least one protein, or at least one RNA, such as, for example, an antisense RNA that is complementary to at least a portion of an mRNA. Typically such a nucleotide construct is comprised of a coding sequence for a protein or an RNA operably linked to 5' and 3' transcriptional regulatory regions. Alternatively, it is also recognized that the methods of the invention may employ a nucleotide construct that is not capable of directing, in a transformed plant, the expression of a protein or an RNA.

In addition, it is recognized that methods of the present invention do not depend on the incorporation into the genome of the entire nucleotide construct comprising a P-glycoprotein nucleotide sequence, only that the plant or cell thereof is altered as a result of the introduction of the nucleotide construct into a cell. In one embodiment of the invention, the genome may be altered following the introduction of the nucleotide construct into a cell. For example, the nucleotide construct, or any part thereof, may incorporate into the genome of the plant. Alterations to the genome of the present invention include, but are not limited to, additions, deletions, and substitutions of nucleotides in the genome. While the methods of the present invention do not depend on additions, deletions, or substitutions of any particular number of nucleotides, it is recognized that such additions, deletions, or substitutions comprise at least one nucleotide.

The nucleotide constructs of the invention also encompass nucleotide constructs that may be employed in methods for altering or mutating a genomic nucleotide sequence in an organism, including, but not limited to, chimeric vectors, chimeric mutational

vectors, chimeric repair vectors, mixed-duplex oligonucleotides, self-complementary chimeric oligonucleotides, and recombinogenic oligonucleobases. Such nucleotide constructs and methods of use, such as, for example, chimeraplasty, are known in the art. Chimeraplasty involves the use of such nucleotide constructs to introduce site-specific changes into the sequence of genomic DNA within an organism. See, U.S. Patent Nos. 5,565,350; 5,731,181; 5,756,325; 5,760,012; 5,795,972; and 5,871,984; all of which are herein incorporated by reference. See also, WO 98/49350, WO 99/07865, WO 99/25821, and Beetham *et al.* (1999) *Proc. Natl. Acad. Sci. USA* 96:8774-8778; herein incorporated by reference.

10 The invention encompasses the use of methods, such as, for example, chimeraplasty to alter P-glycoprotein genes in plants. Such alterations include, for example, changes in the coding sequence that alter the amino acid sequence of the P-glycoprotein encoded thereby, resulting in a reduction in, or loss of, the function of the P-glycoprotein encoded by that gene.

15 The P-glycoprotein nucleotide sequences of the invention are provided in expression cassettes for expression in the plant of interest. The cassette will include 5'- and 3'-regulatory sequences operably linked to a P-glycoprotein nucleotide sequence of the invention. By "operably linked" is intended a functional linkage between a promoter and a second sequence, wherein the promoter sequence initiates and mediates transcription of the DNA sequence corresponding to the second sequence. Generally, operably linked means that the nucleic acid sequences being linked are contiguous and, where necessary to join two protein coding regions, contiguous and in the same reading frame. The cassette may additionally contain at least one additional gene to be cotransformed into the organism. Alternatively, the additional gene(s) can be provided on multiple expression cassettes.

Such an expression cassette is provided with a plurality of restriction sites for insertion of the P-glycoprotein nucleotide sequence to be under the transcriptional regulation of the regulatory regions. The expression cassette may additionally contain selectable marker genes.

30 The expression cassette will include in the 5'-3' direction of transcription, a transcriptional and translational initiation region, a P-glycoprotein nucleotide sequence of

the invention, and a transcriptional and translational termination region functional in plants. The transcriptional initiation region, the promoter, may be native or analogous or foreign or heterologous to the plant host. Additionally, the promoter may be the natural sequence or alternatively a synthetic sequence. By "foreign" is intended that the transcriptional initiation region is not found in the native plant into which the transcriptional initiation region is introduced.

While it may be preferable to express the sequences using heterologous promoters, the native promoter sequences may be used. Such constructs would change expression levels of a P-glycoprotein in the plant or plant cell. Thus, the phenotype of the plant or plant cell is altered.

The termination region may be native with the transcriptional initiation region, may be native with the operably linked DNA sequence of interest, or may be derived from another source. Convenient termination regions are available from the Ti-plasmid of *A. tumefaciens*, such as the octopine synthase and nopaline synthase termination regions. See also Guerineau *et al.* (1991) *Mol. Gen. Genet.* 262:141-144; Proudfoot (1991) *Cell* 64:671-674; Sanfacon *et al.* (1991) *Genes Dev.* 5:141-149; Mogen *et al.* (1990) *Plant Cell* 2:1261-1272; Munroe *et al.* (1990) *Gene* 91:151-158; Ballas *et al.* (1989) *Nucleic Acids Res.* 17:7891-7903; and Joshi *et al.* (1987) *Nucleic Acid Res.* 15:9627-9639.

Where appropriate, the gene(s) may be optimized for increased expression in the transformed plant. That is, the genes can be synthesized using plant-preferred codons for improved expression. Methods are available in the art for synthesizing plant-preferred genes. See, for example, U.S. Patent Nos. 5,380,831, and 5,436,391, and Murray *et al.* (1989) *Nucleic Acids Res.* 17:477-498, herein incorporated by reference.

Additional sequence modifications are known to enhance gene expression in a cellular host. These include elimination of sequences encoding spurious polyadenylation signals, exon-intron splice site signals, transposon-like repeats, and other such well-characterized sequences that may be deleterious to gene expression. The G-C content of the sequence may be adjusted to levels average for a given cellular host, as calculated by reference to known genes expressed in the host cell. When possible, the sequence is modified to avoid predicted hairpin secondary mRNA structures.

U.S. PAT. & TM. OFFICE

The expression cassettes may additionally contain 5'-leader sequences in the expression cassette construct. Such leader sequences can act to enhance translation. Translation leaders are known in the art and include: picornavirus leaders, for example, EMCV leader (Encephalomyocarditis 5'-noncoding region) (Elroy-Stein *et al.* (1989) *PNAS USA* 86:6126-6130); potyvirus leaders, for example, TEV leader (Tobacco Etch Virus) (Allison *et al.* (1986); MDMV leader (Maize Dwarf Mosaic Virus); *Virology* 154:9-20), and human immunoglobulin heavy-chain binding protein (BiP), (Macejak *et al.* (1991) *Nature* 353:90-94); untranslated leader from the coat protein mRNA of alfalfa mosaic virus (AMV RNA 4) (Jobling *et al.* (1987) *Nature* 325:622-625); tobacco mosaic virus leader (TMV) (Gallie *et al.* (1989) in *Molecular Biology of RNA*, ed. Cech (Liss, New York), pp. 237-256); and maize chlorotic mottle virus leader (MCMV) (Lommel *et al.* (1991) *Virology* 81:382-385). See also, Della-Cioppa *et al.* (1987) *Plant Physiol.* 84:965-968. Other methods known to enhance translation can also be utilized, for example, introns, and the like.

15 In preparing the expression cassette, the various DNA fragments may be manipulated, so as to provide for the DNA sequences in the proper orientation and, as appropriate, in the proper reading frame. Toward this end, adapters or linkers may be employed to join the DNA fragments or other manipulations may be involved to provide for convenient restriction sites, removal of superfluous DNA, removal of restriction sites, 20 or the like. For this purpose, *in vitro* mutagenesis, primer repair, restriction, annealing, resubstitutions, e.g., transitions and transversions, may be involved.

It is recognized that with the nucleotide sequences of the invention, antisense constructions, complementary to at least a portion of the messenger RNA (mRNA) for the P-glycoprotein gene sequences can be constructed. Antisense nucleotides are 25 constructed to hybridize with the corresponding mRNA. Modifications of the antisense sequences may be made as long as the sequences hybridize to and interfere with expression of the corresponding mRNA. In this manner, antisense constructions having 70%, preferably 80%, more preferably 85% sequence identity to the corresponding target sequences may be used. Furthermore, portions of the antisense nucleotides may be used 30 to disrupt the expression of the target gene. Generally, sequences of at least 50 nucleotides, 100 nucleotides, 200 nucleotides, or greater may be used.

5 The nucleotide sequences of the present invention may also be used in the sense orientation to suppress the expression of endogenous genes in plants. Methods for suppressing gene expression in plants using nucleotide sequences in the sense orientation, also known as cosuppression methods, are known in the art. The methods generally involve transforming plants with a nucleotide construct comprising a promoter that drives expression in a plant operably linked to at least a portion of a nucleotide sequence that corresponds to the transcript of the endogenous gene. Typically, such a nucleotide sequence has substantial sequence identity to the sequence of the transcript of the endogenous gene, preferably greater than about 65% sequence identity, more preferably greater than about 85% sequence identity, most preferably greater than about 95% sequence identity. See, U.S. Patent Nos. 5,283,184 and 5,034,323; herein incorporated by reference.

15 Generally, the expression cassette will comprise a selectable marker gene for the selection of transformed cells. Selectable marker genes are utilized for the selection of transformed cells or tissues. Marker genes include genes encoding antibiotic resistance, such as those encoding neomycin phosphotransferase II (NEO) and hygromycin phosphotransferase (HPT), as well as genes conferring resistance to herbicidal compounds, such as glufosinate ammonium, bromoxynil, imidazolinones, and 2,4-dichlorophenoxyacetate (2,4-D). See generally, Yarranton (1992) *Curr. Opin. Biotech.* 3:506-511; Christopherson *et al.* (1992) *Proc. Natl. Acad. Sci. USA* 89:6314-6318; Yao *et al.* (1992) *Cell* 71:63-72; Reznikoff (1992) *Mol. Microbiol.* 6:2419-2422; Barkley *et al.* (1980) in *The Operon*, pp. 177-220; Hu *et al.* (1987) *Cell* 48:555-566; Brown *et al.* (1987) *Cell* 49:603-612; Figge *et al.* (1988) *Cell* 52:713-722; Deuschle *et al.* (1989) *Proc. Natl. Acad. Sci. USA* 86:5400-5404; Fuerst *et al.* (1989) *Proc. Natl. Acad. Sci. USA* 86:2549-2553; Deuschle *et al.* (1990) *Science* 248:480-483; Gossen (1993) Ph.D. Thesis, University of Heidelberg; Reines *et al.* (1993) *Proc. Natl. Acad. Sci. USA* 90:1917-1921; Labow *et al.* (1990) *Mol. Cell. Biol.* 10:3343-3356; Zambretti *et al.* (1992) *Proc. Natl. Acad. Sci. USA* 89:3952-3956; Baim *et al.* (1991) *Proc. Natl. Acad. Sci. USA* 88:5072-5076; Wyborski *et al.* (1991) *Nucleic Acids Res.* 19:4647-4653; Hillenand-Wissman (1989) *Topics Mol. Struc. Biol.* 10:143-162; Degenkolb *et al.* (1991) *Antimicrob. Agents Chemother.* 35:1591-1595; Kleinschmidt *et al.* (1988) *Biochemistry* 27:1094-1104; Bonin

(1993) Ph.D. Thesis, University of Heidelberg; Gossen *et al.* (1992) *Proc. Natl. Acad. Sci. USA* 89:5547-5551; Oliva *et al.* (1992) *Antimicrob. Agents Chemother.* 36:913-919; Hlavka *et al.* (1985) *Handbook of Experimental Pharmacology*, Vol. 78 (Springer-Verlag, Berlin); Gill *et al.* (1988) *Nature* 334:721-724. Such disclosures are herein
5 incorporated by reference.

The above list of selectable marker genes is not meant to be limiting. Any selectable marker gene can be used in the present invention.

A number of promoters can be used in the practice of the invention. The promoters may be selected based on the desired timing, localization and level of
10 expression of the P-glycoprotein genes in a plant. Constitutive, tissue-preferred, pathogen-inducible, wound-inducible and chemically regulatable promoters can be used in the practice of the invention.

Such constitutive promoters include, for example, the core promoter of the Rsyn7 promoter and other constitutive promoters disclosed in WO 99/43838 and U.S. Patent
15 No. 6,072,050; the core CaMV 35S promoter (Odell *et al.* (1985) *Nature* 313:810-812); rice actin (McElroy *et al.* (1990) *Plant Cell* 2:163-171); ubiquitin (Christensen *et al.* (1989) *Plant Mol. Biol.* 12:619-632 and Christensen *et al.* (1992) *Plant Mol. Biol.* 18:675-689); pEMU (Last *et al.* (1991) *Theor. Appl. Genet.* 81:581-588); MAS (Velten *et al.* (1984) *EMBO J.* 3:2723-2730); ALS promoter (U.S. Application Serial No.
20 08/409,297), and the like. Other constitutive promoters include, for example, U.S. Patent Nos. 5,608,149; 5,608,144; 5,604,121; 5,569,597; 5,466,785; 5,399,680; 5,268,463; and 5,608,142.

Tissue-preferred promoters can be utilized to target enhanced P-glycoprotein expression within a particular plant tissue. Tissue-preferred promoters include
25 Yamamoto *et al.* (1997) *Plant J.* 12(2):255-265; Kawamata *et al.* (1997) *Plant Cell Physiol.* 38(7):792-803; Hansen *et al.* (1997) *Mol. Gen Genet.* 254(3):337-343; Russell *et al.* (1997) *Transgenic Res.* 6(2):157-168; Rinehart *et al.* (1996) *Plant Physiol.* 112(3):1331-1341; Van Camp *et al.* (1996) *Plant Physiol.* 112(2):525-535; Canevascini *et al.* (1996) *Plant Physiol.* 112(2):513-524; Yamamoto *et al.* (1994) *Plant Cell Physiol.*
30 35(5):773-778; Lam (1994) *Results Probl. Cell Differ.* 20:181-196; Orozco *et al.* (1993) *Plant Mol Biol.* 23(6):1129-1138; Matsuoka *et al.* (1993) *Proc Natl. Acad. Sci. USA*

90(20):9586-9590; and Guevara-Garcia *et al.* (1993) *Plant J.* 4(3):495-505. Such promoters can be modified, if necessary, for weak expression.

Leaf-preferred promoters include, Yamamoto *et al.* (1997) *Plant J.* 12(2):255-265; Kawamata *et al.* (1997) *Plant Cell Physiol.* 38(7):792-803; Hansen *et al.* (1997) *Mol. Gen. Genet.* 254(3):337-343; Russell *et al.* (1997) *Transgenic Res.* 6(2):157-168; Rinehart *et al.* (1996) *Plant Physiol.* 112(3):1331-1341; Van Camp *et al.* (1996) *Plant Physiol.* 112(2):525-535; Canevascini *et al.* (1996) *Plant Physiol.* 112(2):513-524; Yamamoto *et al.* (1994) *Plant Cell Physiol.* 35(5):773-778; Lam (1994) *Results Probl. Cell Differ.* 20:181-196; Orozco *et al.* (1993) *Plant Mol. Biol.* 23(6):1129-1138; Matsuoka *et al.* (1993) *Proc. Natl. Acad. Sci. USA* 90(20):9586-9590; and Guevara-Garcia *et al.* (1993) *Plant J.* 4(3):495-505.

Root-preferred promoters are known and can be selected from the many available from the literature or isolated de novo from various compatible species. See, for example, Hire *et al.* (1992) *Plant Mol. Biol.* 20(2): 207-218 (soybean root-preferred glutamine synthetase gene); Keller and Baumgartner (1991) *Plant Cell* 3(10):1051-1061 (root-preferred control element in the GRP 1.8 gene of French bean); Sanger *et al.* (1990) *Plant Mol. Biol.* 14(3):433-443 (root-preferred promoter of the mannopine synthase (MAS) gene of *Agrobacterium tumefaciens*); and Miao *et al.* (1991) *Plant Cell* 3(1):11-22 (full-length cDNA clone encoding cytosolic glutamine synthetase (GS), which is expressed in roots and root nodules of soybean). See also Bogusz *et al.* (1990) *Plant Cell* 2(7):633-641, where two root-preferred promoters isolated from hemoglobin genes from the nitrogen-fixing nonlegume *Parasponia andersonii* and the related non-nitrogen-fixing nonlegume *Trema tomentosa* are described. The promoters of these genes were linked to a β -glucuronidase reporter gene and introduced into both the nonlegume *Nicotiana tabacum* and the legume *Lotus corniculatus*, and in both instances root-preferred promoter activity was preserved. Leach and Aoyagi (1991) describe their analysis of the promoters of the highly expressed rolC and rolD root-inducing genes of *Agrobacterium rhizogenes* (see *Plant Science* (Limerick) 79(1):69-76). They concluded that enhancer and tissue-preferred DNA determinants are dissociated in those promoters. Teeri *et al.* (1989) used gene fusion to lacZ to show that the *Agrobacterium* T-DNA gene encoding octopine synthase is especially active in the epidermis of the root tip and that the TR2'

gene is root preferred in the intact plant and stimulated by wounding in leaf tissue, an especially desirable combination of characteristics for use with an insecticidal or larvicidal gene (see *EMBO J.* 8(2):343-350). The TR1' gene, fused to *nptII* (neomycin phosphotransferase II) showed similar characteristics. Additional root-preferred promoters include the VfENOD-GRP3 gene promoter (Kuster *et al.* (1995) *Plant Mol. Biol.* 29(4):759-772); and rolB promoter (Capano *et al.* (1994) *Plant Mol. Biol.* 25(4):681-691. See also U.S. Patent Nos. 5,837,876; 5,750,386; 5,633,363; 5,459,252; 5,401,836; 5,110,732; and 5,023,179.

Generally, it will be beneficial to express the gene from an inducible promoter, particularly from a pathogen-inducible promoter. Such promoters include those from pathogenesis-related proteins (PR proteins), which are induced following infection by a pathogen; e.g., PR proteins, SAR proteins, beta-1,3-glucanase, chitinase, etc. See, for example, Redolfi *et al.* (1983) *Neth. J. Plant Pathol.* 89:245-254; Uknes *et al.* (1992) *Plant Cell* 4:645-656; and Van Loon (1985) *Plant Mol. Virol.* 4:111-116. See also the copending applications entitled "Inducible Maize Promoters", U.S. Application Serial No. 60/076,100, filed February 26, 1998, and U.S. Application Serial No. 60/079,648, filed March 27, 1998, both of which are herein incorporated by reference.

Of interest are promoters that are expressed locally at or near the site of pathogen infection. See, for example, Marineau *et al.* (1987) *Plant Mol. Biol.* 9:335-342; Matton *et al.* (1989) *Molecular Plant-Microbe Interactions* 2:325-331; Somsisch *et al.* (1986) *Proc. Natl. Acad. Sci. USA* 83:2427-2430; Somsisch *et al.* (1988) *Mol. Gen. Genet.* 2:93-98; and Yang (1996) *Proc. Natl. Acad. Sci. USA* 93:14972-14977. See also, Chen *et al.* (1996) *Plant J.* 10:955-966; Zhang *et al.* (1994) *Proc. Natl. Acad. Sci. USA* 91:2507-2511; Warner *et al.* (1993) *Plant J.* 3:191-201; Siebertz *et al.* (1989) *Plant Cell* 1:961-968; U.S. Patent No. 5,750,386 (nematode-inducible); and the references cited therein. Of particular interest is the inducible promoter for the maize PRms gene, whose expression is induced by the pathogen *Fusarium moniliforme* (see, for example, Cordero *et al.* (1992) *Physiol. Mol. Plant Path.* 41:189-200).

Additionally, as pathogens find entry into plants through wounds or insect damage, a wound-inducible promoter may be used in the constructions of the invention. Such wound-inducible promoters include potato proteinase inhibitor (pin II) gene (Ryan

(1990) *Ann. Rev. Phytopath.* 28:425-449; Duan *et al.* (1996) *Nature Biotechnology* 14:494-498; wun1 and wun2, US Patent No. 5,428,148; win1 and win2 (Stanford *et al.* (1989) *Mol. Gen. Genet.* 215:200-208); systemin (McGurl *et al.* (1992) *Science* 225:1570-1573); WIP1 (Rohmeier *et al.* (1993) *Plant Mol. Biol.* 22:783-792; Eckelkamp *et al.* (1993) *FEBS Letters* 323:73-76); MPI gene (Corderok *et al.* (1994) *Plant J.* 6(2):141-150); and the like, herein incorporated by reference.

Chemically regulated promoters can be used to modulate the expression of a gene in a plant through the application of an exogenous chemical regulator. Depending upon the objective, the promoter may be a chemical-inducible promoter, where application of the chemical induces gene expression, or a chemical-repressible promoter, where application of the chemical represses gene expression. Chemically inducible promoters are known in the art and include, but are not limited to, the maize In2-2 promoter, which is activated by benzenesulfonamide herbicide safeners, the maize GST promoter, which is activated by hydrophobic electrophilic compounds that are used as pre-emergent herbicides, and the tobacco PR-1a promoter, which is activated by salicylic acid. Other chemically regulated promoters of interest include steroid-responsive promoters (see, for example, the glucocorticoid-inducible promoter in Schena *et al.* (1991) *Proc. Natl. Acad. Sci. USA* 88:10421-10425 and McNellis *et al.* (1998) *Plant J.* 14(2):247-257) and tetracycline-inducible and tetracycline-repressible promoters (see, for example, Gatz *et al.* (1991) *Mol. Gen. Genet.* 227:229-237, and U.S. Patent Nos. 5,814,618 and 5,789,156), herein incorporated by reference.

Transformation protocols as well as protocols for introducing nucleotide sequences into plants may vary depending on the type of plant or plant cell, i.e., monocot or dicot, targeted for transformation. Suitable methods of introducing nucleotide sequences into plant cells and subsequent insertion into the plant genome include microinjection (Crossway *et al.* (1986) *Biotechniques* 4:320-334), electroporation (Riggs *et al.* (1986) *Proc. Natl. Acad. Sci. USA* 83:5602-5606, *Agrobacterium*-mediated transformation (Townsend *et al.*, U.S. Patent No. 5,563,055; Zhao *et al.*, U.S. Patent No. 5,981,840), direct gene transfer (Paszkowski *et al.* (1984) *EMBO J.* 3:2717-2722), and ballistic particle acceleration (see, for example, Sanford *et al.*, U.S. Patent No. 4,945,050; Tomes *et al.* (1995) "Direct DNA Transfer into Intact Plant Cells via Microprojectile

Bombardment,” in *Plant Cell, Tissue, and Organ Culture: Fundamental Methods*, ed. Gamborg and Phillips (Springer-Verlag, Berlin); and McCabe *et al.* (1988) *Biotechnology* 6:923-926). Also see Weissinger *et al.* (1988) *Ann. Rev. Genet.* 22:421-477; Sanford *et al.* (1987) *Particulate Science and Technology* 5:27-37 (onion); Christou *et al.* (1988) *Plant Physiol.* 87:671-674 (soybean); McCabe *et al.* (1988) *Bio/Technology* 6:923-926 (soybean); Finer and McMullen (1991) *In Vitro Cell Dev. Biol.* 27P:175-182 (soybean); Singh *et al.* (1998) *Theor. Appl. Genet.* 96:319-324 (soybean); Datta *et al.* (1990) *Biotechnology* 8:736-740 (rice); Klein *et al.* (1988) *Proc. Natl. Acad. Sci. USA* 85:4305-4309 (maize); Klein *et al.* (1988) *Biotechnology* 6:559-563 (maize); Tomes, U.S. Patent No. 5,240,855; Buising *et al.*, U.S. Patent Nos. 5,322,783 and 5,324,646; Tomes *et al.* (1995) “Direct DNA Transfer into Intact Plant Cells via Microprojectile Bombardment,” in *Plant Cell, Tissue, and Organ Culture: Fundamental Methods*, ed. Gamborg (Springer-Verlag, Berlin) (maize); Klein *et al.* (1988) *Plant Physiol.* 91:440-444 (maize); Fromm *et al.* (1990) *Biotechnology* 8:833-839 (maize); Hooykaas-Van Slogteren *et al.* (1984) *Nature (London)* 311:763-764; Bytebier *et al.* (1987) *Proc. Natl. Acad. Sci. USA* 84:5345-5349 (Liliaceae); De Wet *et al.* (1985) in *The Experimental Manipulation of Ovule Tissues*, ed. Chapman *et al.* (Longman, New York), pp. 197-209 (pollen); Kaeppler *et al.* (1990) *Plant Cell Reports* 9:415-418 and Kaeppler *et al.* (1992) *Theor. Appl. Genet.* 84:560-566 (whisker-mediated transformation); D'Halluin *et al.* (1992) *Plant Cell* 4:1495-1505 (electroporation); Li *et al.* (1993) *Plant Cell Reports* 12:250-255 and Christou and Ford (1995) *Annals of Botany* 75:407-413 (rice); Osjoda *et al.* (1996) *Nature Biotechnology* 14:745-750 (maize via *Agrobacterium tumefaciens*); all of which are herein incorporated by reference.

Alternatively, the nucleotide sequences of the invention can be introduced into an organism and allowed to undergo recombination with homologous regions of the organism's genome. Such homologous recombination approaches are well known to those of ordinary skill in the art and can be used to stably incorporate sequences of the invention into an organism. Further, such strategies can be used to introduce “knockout mutations” into a specific gene of an organism that shares substantial homology to the sequences of the invention. A knockout mutation is any mutation in the sequence of a gene that eliminates or substantially reduces the function or the level of the product

encoded by the gene. Methods involving transformation of an organism followed by homologous recombination to stably integrate the sequences of the invention into the genome organism are encompassed by the invention. The invention is particularly directed to methods where sequences of the invention are utilized to alter the growth of an organism. Such methods encompass use of the sequences of the invention to interfere with the function or synthesis of a P-glycoprotein that controls growth of an organism.

The cells that have been transformed may be grown into plants in accordance with conventional ways. See, for example, McCormick *et al.* (1986) *Plant Cell Reports* 5:81-84. These plants may then be grown, and either pollinated with the same transformed strain or different strains, and the resulting hybrid having constitutive expression of the desired phenotypic characteristic identified. Two or more generations may be grown to ensure that constitutive expression of the desired phenotypic characteristic is stably maintained and inherited and then seeds harvested to ensure constitutive expression of the desired phenotypic characteristic has been achieved.

The present invention may be used for transformation of any plant species, including, but not limited to, corn (*Zea mays*), *Brassica* sp. (e.g., *B. napus*, *B. rapa*, *B. juncea*), particularly those *Brassica* species useful as sources of seed oil, alfalfa (*Medicago sativa*), rice (*Oryza sativa*), rye (*Secale cereale*), sorghum (*Sorghum bicolor*, *Sorghum vulgare*), millet (e.g., pearl millet (*Pennisetum glaucum*), proso millet (*Panicum miliaceum*), foxtail millet (*Setaria italica*), finger millet (*Eleusine coracana*)), sunflower (*Helianthus annuus*), safflower (*Carthamus tinctorius*), wheat (*Triticum aestivum*), soybean (*Glycine max*), tobacco (*Nicotiana tabacum*), potato (*Solanum tuberosum*), peanuts (*Arachis hypogaea*), cotton (*Gossypium barbadense*, *Gossypium hirsutum*), sweet potato (*Ipomoea batatas*), cassava (*Manihot esculenta*), coffee (*Coffea* spp.), coconut (*Cocos nucifera*), pineapple (*Ananas comosus*), citrus trees (*Citrus* spp.), cocoa (*Theobroma cacao*), tea (*Camellia sinensis*), banana (*Musa* spp.), avocado (*Persea americana*), fig (*Ficus casica*), guava (*Psidium guajava*), mango (*Mangifera indica*), olive (*Olea europaea*), papaya (*Carica papaya*), cashew (*Anacardium occidentale*), macadamia (*Macadamia integrifolia*), almond (*Prunus amygdalus*), sugar beets (*Beta vulgaris*), sugarcane (*Saccharum* spp.), oats, barley, vegetables, ornamentals, and conifers.

Vegetables include tomatoes (*Lycopersicon esculentum*), lettuce (e.g., *Lactuca sativa*), green beans (*Phaseolus vulgaris*), lima beans (*Phaseolus limensis*), peas (*Lathyrus* spp.), and members of the genus *Cucumis* such as cucumber (*C. sativus*), cantaloupe (*C. cantalupensis*), and musk melon (*C. melo*). Ornamentals include azalea
5 (*Rhododendron* spp.), hydrangea (*Macrophylla hydrangea*), hibiscus (*Hibiscus rosasanensis*), roses (*Rosa* spp.), tulips (*Tulipa* spp.), daffodils (*Narcissus* spp.), petunias (*Petunia hybrida*), carnation (*Dianthus caryophyllus*), poinsettia (*Euphorbia pulcherrima*), and chrysanthemum. Conifers that may be employed in practicing the present invention include, for example, pines such as loblolly pine (*Pinus taeda*), slash
10 pine (*Pinus elliotii*), ponderosa pine (*Pinus ponderosa*), lodgepole pine (*Pinus contorta*), and Monterey pine (*Pinus radiata*); Douglas-fir (*Pseudotsuga menziesii*); Western hemlock (*Tsuga canadensis*); Sitka spruce (*Picea glauca*); redwood (*Sequoia sempervirens*); true firs such as silver fir (*Abies amabilis*) and balsam fir (*Abies balsamea*); and cedars such as Western red cedar (*Thuja plicata*) and Alaska yellow-
15 cedar (*Chamaecyparis nootkatensis*). Preferably, plants of the present invention are crop plants (for example, rice, corn, alfalfa, sunflower, *Brassica*, soybean, cotton, safflower, peanut, sorghum, wheat, millet, tobacco, etc.), more preferably corn, rice and sorghum plants.

The invention is drawn to compositions and methods for increasing the resistance
20 of a plant to a pathogen. Accordingly, the compositions and methods are also useful in protecting plants against fungal pathogens, viruses, nematodes, insects, acarids and the like.

By “disease resistance” is intended that the plants avoid the disease symptoms that are the outcome of plant-pathogen interactions. That is, pathogens are prevented
25 from causing plant diseases and the associated disease symptoms, or alternatively, the disease symptoms caused by the pathogen is minimized or lessened. The methods of the invention can be utilized to protect plants from disease, particularly those diseases that are caused by plant pathogens.

By “antipathogenic compositions” is intended that the compositions of the
30 invention have antipathogenic activity and thus are capable of suppressing, controlling, and/or killing the invading pathogenic organism. An antipathogenic composition of the

invention will reduce the disease symptoms resulting from pathogen challenge by at least about 5% to about 50%, at least about 10% to about 60%, at least about 30% to about 70%, at least about 40% to about 80%, or at least about 50% to about 90% or greater. Hence, the methods of the invention can be utilized to protect plants from disease,
5 particularly those diseases that are caused by plant pathogens.

Assays that measure antipathogenic activity are commonly known in the art, as are methods to quantitate disease resistance in plants following pathogen infection. See, for example, U.S. Patent No. 5,614,395, herein incorporated by reference. Such techniques include, measuring over time, the average lesion diameter, the pathogen
10 biomass, and the overall percentage of decayed plant tissues. For example, a plant either expressing an antipathogenic polypeptide or having an antipathogenic composition applied to its surface shows a decrease in tissue necrosis (*i.e.*, lesion diameter) or a decrease in plant death following pathogen challenge when compared to a control plant that was not exposed to the antipathogenic composition. Alternatively, antipathogenic
15 activity can be measured by a decrease in pathogen biomass. For example, a plant expressing an antipathogenic polypeptide or exposed to an antipathogenic composition is challenged with a pathogen of interest. Over time, tissue samples from the pathogen-inoculated tissues are obtained and RNA is extracted. The percent of a specific pathogen RNA transcript relative to the level of a plant specific transcript allows the level of
20 pathogen biomass to be determined. See, for example, Thomma *et al.* (1998) *Plant Biology* 95:15107-15111, herein incorporated by reference.

Furthermore, *in vitro* antipathogenic assays include, for example, the addition of varying concentrations of the antipathogenic composition to paper disks and placing the disks on agar containing a suspension of the pathogen of interest. Following incubation,
25 clear inhibition zones develop around the discs that contain an effective concentration of the antipathogenic polypeptide (Liu *et al.* (1994) *Plant Biology* 91:1888-1892, herein incorporated by reference). Additionally, microspectrophotometrical analysis can be used to measure the *in vitro* antipathogenic properties of a composition (Hu *et al.* (1997) *Plant Mol. Biol.* 34:949-959 and Cammue *et al.* (1992) *J. Biol. Chem.* 267: 2228-2233,
30 both of which are herein incorporated by reference).

Pathogens of the invention include, but are not limited to, viruses or viroids, bacteria, insects, nematodes, fungi, and the like. Viruses include any plant virus, for example, tobacco or cucumber mosaic virus, ringspot virus, necrosis virus, maize dwarf mosaic virus, etc. Specific fungal and viral pathogens for the major crops include:

- 5 Soybeans: *Phytophthora megasperma* fsp. *glycinea*, *Macrophomina phaseolina*, *Rhizoctonia solani*, *Sclerotinia sclerotiorum*, *Fusarium oxysporum*, *Diaporthe phaseolorum* var. *sojae* (*Phomopsis sojae*), *Diaporthe phaseolorum* var. *caulivora*, *Sclerotium rolfsii*, *Cercospora kikuchii*, *Cercospora sojae*, *Peronospora manshurica*, *Colletotrichum dematium* (*Colletotrichum truncatum*), *Corynespora cassiicola*, *Septoria*
- 10 *glycines*, *Phyllosticta sojicola*, *Alternaria alternata*, *Pseudomonas syringae* p.v. *glycinea*, *Xanthomonas campestris* p.v. *phaseoli*, *Microsphaera diffusa*, *Fusarium semitectum*, *Phialophora gregata*, Soybean mosaic virus, *Glomerella glycines*, Tobacco Ring spot virus, Tobacco Streak virus, *Phakopsora pachyrhizi*, *Pythium aphanidermatum*, *Pythium ultimum*, *Pythium debaryanum*, Tomato spotted wilt virus, *Heterodera glycines* *Fusarium*
- 15 *solani*; Canola: *Albugo candida*, *Alternaria brassicae*, *Leptosphaeria maculans*, *Rhizoctonia solani*, *Sclerotinia sclerotiorum*, *Mycosphaerella brassicicola*, *Pythium ultimum*, *Peronospora parasitica*, *Fusarium roseum*, *Alternaria alternata*; Alfalfa: *Clavibacter michiganense* subsp. *insidiosum*, *Pythium ultimum*, *Pythium irregulare*, *Pythium splendens*, *Pythium debaryanum*, *Pythium aphanidermatum*, *Phytophthora*
- 20 *megasperma*, *Peronospora trifoliorum*, *Phoma medicaginis* var. *medicaginis*, *Cercospora medicaginis*, *Pseudopeziza medicaginis*, *Leptotrochila medicaginis*, *Fusarium*, *Xanthomonas campestris* p.v. *alfalfae*, *Aphanomyces euteiches*, *Stemphylium herbarum*, *Stemphylium alfalfae*; Wheat: *Pseudomonas syringae* p.v. *atrofaciens*, *Urocystis agropyri*, *Xanthomonas campestris* p.v. *translucens*, *Pseudomonas syringae* p.v.
- 25 *syringae*, *Alternaria alternata*, *Cladosporium herbarum*, *Fusarium graminearum*, *Fusarium avenaceum*, *Fusarium culmorum*, *Ustilago tritici*, *Ascochyta tritici*, *Cephalosporium gramineum*, *Collotetrichum graminicola*, *Erysiphe graminis* f.sp. *tritici*, *Puccinia graminis* f.sp. *tritici*, *Puccinia recondita* f.sp. *tritici*, *Puccinia striiformis*, *Pyrenophora tritici-repentis*, *Septoria nodorum*, *Septoria tritici*, *Septoria avenae*,
- 30 *Pseudocercospora herpotrichoides*, *Rhizoctonia solani*, *Rhizoctonia cerealis*, *Gaeumannomyces graminis* var. *tritici*, *Pythium aphanidermatum*, *Pythium*

- arrhenomanes*, *Pythium ultimum*, *Bipolaris sorokiniana*, Barley Yellow Dwarf Virus, Brome Mosaic Virus, Soil Borne Wheat Mosaic Virus, Wheat Streak Mosaic Virus, Wheat Spindle Streak Virus, American Wheat Striate Virus, *Claviceps purpurea*, *Tilletia tritici*, *Tilletia laevis*, *Ustilago tritici*, *Tilletia indica*, *Rhizoctonia solani*, *Pythium*
- 5 *arrhenomannes*, *Pythium graminicola*, *Pythium aphanidermatum*, High Plains Virus, European wheat striate virus; Sunflower: *Plasmophora halstedii*, *Sclerotinia sclerotiorum*, Aster Yellows, *Septoria helianthi*, *Phomopsis helianthi*, *Alternaria helianthi*, *Alternaria zinniae*, *Botrytis cinerea*, *Phoma macdonaldii*, *Macrophomina phaseolina*, *Erysiphe cichoracearum*, *Rhizopus oryzae*, *Rhizopus arrhizus*, *Rhizopus*
- 10 *stolonifer*, *Puccinia helianthi*, *Verticillium dahliae*, *Erwinia carotovorum* pv. *carotovora*, *Cephalosporium acremonium*, *Phytophthora cryptogea*, *Albugo tragopogonis*; Corn: *Fusarium moniliforme* var. *subglutinans*, *Erwinia stewartii*, *Fusarium moniliforme*, *Gibberella zeae* (*Fusarium graminearum*), *Stenocarpella maydi* (*Diplodia maydis*), *Pythium irregulare*, *Pythium debaryanum*, *Pythium graminicola*, *Pythium splendens*,
- 15 *Pythium ultimum*, *Pythium aphanidermatum*, *Aspergillus flavus*, *Bipolaris maydis* O, T (*Cochliobolus heterostrophus*), *Helminthosporium carbonum* I, II & III (*Cochliobolus carbonum*), *Exserohilum turcicum* I, II & III, *Helminthosporium pedicellatum*, *Physoderma maydis*, *Phyllosticta maydis*, *Kabatiella-maydis*, *Cercospora sorghi*, *Ustilago maydis*, *Puccinia sorghi*, *Puccinia polysora*, *Macrophomina phaseolina*,
- 20 *Penicillium oxalicum*, *Nigrospora oryzae*, *Cladosporium herbarum*, *Curvularia lunata*, *Curvularia inaequalis*, *Curvularia pallescens*, *Clavibacter michiganense* subsp. *nebraskense*, *Trichoderma viride*, Maize Dwarf Mosaic Virus A & B, Wheat Streak Mosaic Virus, Maize Chlorotic Dwarf Virus, *Claviceps sorghi*, *Pseudonomas avenae*, *Erwinia chrysanthemi* pv. *zea*, *Erwinia carotovora*, Corn stunt Spiroplasma, *Diplodia*
- 25 *macrospora*, *Sclerophthora macrospora*, *Peronosclerospora sorghi*, *Peronosclerospora philippinensis*, *Peronosclerospora maydis*, *Peronosclerospora sacchari*, *Sphacelotheca reiliana*, *Physopella zeae*, *Cephalosporium maydis*, *Cephalosporium acremonium*, Maize Chlorotic Mottle Virus, High Plains Virus, Maize Mosaic Virus, Maize Rayado Fino Virus, Maize Streak Virus, Maize Stripe Virus, Maize Rough Dwarf Virus; Sorghum:
- 30 *Exserohilum turcicum*, *Colletotrichum graminicola* (*Glomerella graminicola*), *Cercospora sorghi*, *Gloeocercospora sorghi*, *Ascochyta sorghina*, *Pseudomonas syringae*

p.v. *syringae*, *Xanthomonas campestris* p.v. *holcicola*, *Pseudomonas andropogonis*,
Puccinia purpurea, *Macrophomina phaseolina*, *Perconia circinata*, *Fusarium*
moniliforme, *Alternaria alternata*, *Bipolaris sorghicola*, *Helminthosporium sorghicola*,
Curvularia lunata, *Phoma insidiosa*, *Pseudomonas avenae* (*Pseudomonas*
5 *alboprecipitans*), *Ramulispora sorghi*, *Ramulispora sorghicola*, *Phyllachara sacchari*,
Sporisorium reilianum (*Sphacelotheca reiliana*), *Sphacelotheca cruenta*, *Sporisorium*
sorghii, Sugarcane mosaic H, Maize Dwarf Mosaic Virus A & B, *Claviceps sorghi*,
Rhizoctonia solani, *Acremonium strictum*, *Sclerophthora macrospora*,
Peronosclerospora sorghi, *Peronosclerospora philippinensis*, *Sclerospora graminicola*,
10 *Fusarium graminearum*, *Fusarium oxysporum*, *Pythium arrhenomanes*, *Pythium*
graminicola, etc.

Nematodes include parasitic nematodes such as root-knot, cyst, and lesion
nematodes, including *Heterodera* and *Globodera* spp; particularly *Globodera*
rostochiensis and *globodera pailida* (potato cyst nematodes); *Heterodera glycines*
15 (soybean cyst nematode); *Heterodera schachtii* (beet cyst nematode); and *Heterodera*
avenae (cereal cyst nematode).

Insect pests include insects selected from the orders Coleoptera, Diptera,
Hymenoptera, Lepidoptera, Mallophaga, Homoptera, Hemiptera, Orthoptera,
Thysanoptera, Dermaptera, Isoptera, Anoplura, Siphonaptera, Trichoptera, etc.,
20 particularly Coleoptera and Lepidoptera. Insect pests of the invention for the major crops
include: Maize: *Ostrinia nubilalis*, European corn borer; *Agrotis ipsilon*, black cutworm;
Helicoverpa zea, corn earworm; *Spodoptera frugiperda*, fall armyworm; *Diatraea*
grandiosella, southwestern corn borer; *Elasmopalpus lignosellus*, lesser cornstalk borer;
Diatraea saccharalis, sugarcane borer; *Diabrotica virgifera*, western corn rootworm;
25 *Diabrotica longicornis barberi*, northern corn rootworm; *Diabrotica undecimpunctata*
howardi, southern corn rootworm; *Melanotus* spp., wireworms; *Cyclocephala borealis*,
northern masked chafer (white grub); *Cyclocephala immaculata*, southern masked chafer
(white grub); *Popillia japonica*, Japanese beetle; *Chaetocnema pulicaria*, corn flea
beetle; *Sphenophorus maidis*, maize billbug; *Rhopalosiphum maidis*, corn leaf aphid;
30 *Anuraphis maidiradicis*, corn root aphid; *Blissus leucopterus leucopterus*, chinch bug;
Melanoplus femurrubrum, redlegged grasshopper; *Melanoplus sanguinipes*, migratory

- grasshopper; *Hylemya platura*, seedcorn maggot; *Agromyza parvicornis*, corn blot
leafminer; *Anaphothrips obscurus*, grass thrips; *Solenopsis milesta*, thief ant;
Tetranychus urticae, twospotted spider mite; Sorghum: *Chilo partellus*, sorghum borer;
Spodoptera frugiperda, fall armyworm; *Helicoverpa zea*, corn earworm; *Elasmopalpus*
5 *lignosellus*, lesser cornstalk borer; *Feltia subterranea*, granulate cutworm; *Phyllophaga*
crinita, white grub; *Eleodes*, *Conoderus*, and *Aeolus spp.*, wireworms; *Oulema*
melanopus, cereal leaf beetle; *Chaetocnema pulicaria*, corn flea beetle; *Sphenophorus*
maidis, maize billbug; *Rhopalosiphum maidis*; corn leaf aphid; *Sipha flava*, yellow
sugarcane aphid; *Blissus leucopterus leucopterus*, chinch bug; *Contarinia sorghicola*,
10 sorghum midge; *Tetranychus cinnabarinus*, carmine spider mite; *Tetranychus urticae*,
twospotted spider mite; Wheat: *Pseudaletia unipunctata*, army worm; *Spodoptera*
frugiperda, fall armyworm; *Elasmopalpus lignosellus*, lesser cornstalk borer; *Agrotis*
orthogonia, western cutworm; *Elasmopalpus lignosellus*, lesser cornstalk borer; *Oulema*
melanopus, cereal leaf beetle; *Hypera punctata*, clover leaf weevil; *Diabrotica*
15 *undecimpunctata howardi*, southern corn rootworm; Russian wheat aphid; *Schizaphis*
graminum, greenbug; *Macrosiphum avenae*, English grain aphid; *Melanoplus*
femurrubrum, redlegged grasshopper; *Melanoplus differentialis*, differential grasshopper;
Melanoplus sanguinipes, migratory grasshopper; *Mayetiola destructor*, Hessian fly;
Sitodiplosis mosellana, wheat midge; *Meromyza americana*, wheat stem maggot;
20 *Hylemya coarctata*, wheat bulb fly; *Frankliniella fusca*, tobacco thrips; *Cephus cinctus*,
wheat stem sawfly; *Aceria tulipae*, wheat curl mite; Sunflower: *Suleima helianthana*,
sunflower bud moth; *Homoeosoma electellum*, sunflower moth; *zygogramma*
exclamationis, sunflower beetle; *Bothyrus gibbosus*, carrot beetle; *Neolasioptera*
murtfeldtiana, sunflower seed midge; Cotton: *Heliothis virescens*, cotton budworm;
25 *Helicoverpa zea*, cotton bollworm; *Spodoptera exigua*, beet armyworm; *Pectinophora*
gossypiella, pink bollworm; *Anthonomus grandis grandis*, boll weevil; *Aphis gossypii*,
cotton aphid; *Pseudatomoscelis seriatus*, cotton fleahopper; *Trialeurodes abutilonea*,
bandedwinged whitefly; *Lygus lineolaris*, tarnished plant bug; *Melanoplus femurrubrum*,
redlegged grasshopper; *Melanoplus differentialis*, differential grasshopper; *Thrips tabaci*,
30 onion thrips; *Frankliniella fusca*, tobacco thrips; *Tetranychus cinnabarinus*, carmine
spider mite; *Tetranychus urticae*, twospotted spider mite; Rice: *Diatraea saccharalis*,

sugarcane borer; *Spodoptera frugiperda*, fall armyworm; *Helicoverpa zea*, corn earworm; *Colaspis brunnea*, grape colaspis; *Lissorhoptrus oryzophilus*, rice water weevil; *Sitophilus oryzae*, rice weevil; *Nephotettix nigropictus*, rice leafhopper; *Blissus leucopterus leucopterus*, chinch bug; *Acrosternum hilare*, green stink bug; Soybean:
5 *Pseudoplusia includens*, soybean looper; *Anticarsia gemmatilis*, velvetbean caterpillar; *Plathypena scabra*, green cloverworm; *Ostrinia nubilalis*, European corn borer; *Agrotis ipsilon*, black cutworm; *Spodoptera exigua*, beet armyworm; *Heliothis virescens*, cotton budworm; *Helicoverpa zea*, cotton bollworm; *Epilachna varivestis*, Mexican bean beetle; *Myzus persicae*, green peach aphid; *Empoasca fabae*, potato leafhopper; *Acrosternum*
10 *hilare*, green stink bug; *Melanoplus femurrubrum*, redlegged grasshopper; *Melanoplus differentialis*, differential grasshopper; *Hylemya platura*, seedcorn maggot; *Sericothrips variabilis*, soybean thrips; *Thrips tabaci*, onion thrips; *Tetranychus turkestani*, strawberry spider mite; *Tetranychus urticae*, twospotted spider mite; Barley: *Ostrinia nubilalis*, European corn borer; *Agrotis ipsilon*, black cutworm; *Schizaphis graminum*, greenbug;
15 *Blissus leucopterus leucopterus*, chinch bug; *Acrosternum hilare*, green stink bug; *Euschistus servus*, brown stink bug; *Delia platura*, seedcorn maggot; *Mayetiola destructor*, Hessian fly; *Petrobia latens*, brown wheat mite; Oil Seed Rape: *Brevicoryne brassicae*, cabbage aphid; *Phyllotreta cruciferae*, Flea beetle; *Mamestra configurata*, Bertha armyworm; *Plutella xylostella*, Diamond-back moth; *Delia* ssp., Root maggots.
20 The following examples are offered by way of illustration and not by way of limitation.

EXAMPLE 1

Sorghum dwarfing gene, *Dw3*, encodes a P-glycoprotein homologue

25 It is well established that the sorghum dwarfing phenotype conferred by the *dw3* recessive mutation is unstable, although the mechanism responsible for its instability remains unknown. The *dw3* allele, referred to here as the reference allele (*dw3-ref*), reverts back to the wild-type form (conferring a tall phenotype) with a frequency of about 0.4 % to about 1 %. As a result, it is a commonplace to witness a number of tall sorghum
30 plants in a field of *dw3* dwarfs. To determine if there is any relationship between the maize *br2* gene and the sorghum *dw3* gene, leaf samples were collected from 8 dwarf and

8 tall (revertant) plants; these were expected to be true *dw3* isogenics (identical throughout the genome except at the *dw3* locus). The DNA of these samples was extracted, digested with *Pst*I, and subjected to Southern blot analysis using a probe from the maize *Br2* gene. A clear and consistent DNA polymorphism was observed between
5 the tall and dwarf plants, with the restriction fragment from the revertant allele being about 1.0 kb smaller than the *dw3-ref* allele.

Two conclusions were made from this result. First, the sorghum *Dw3* locus is structurally and functionally homologous to the maize *Br2* gene, suggesting that they may turn out to be true orthologs (i.e., derived from the same ancestral gene by vertical
10 descent). Second, since all revertants had the same RFLP pattern, and that the size of the revertant allele was smaller than the mutant allele, the mutable *dw3-ref* allele was probably caused by an insertion. To address the latter interpretation, sorghum DNA in the vicinity of the *Br2*-detected polymorphism was subjected to PCR amplification using two oligonucleotide primers (SEQ ID NOS: 5-6) derived from the nucleotide sequence of
15 the maize *Br2* gene.

PCR products were amplified from genomic DNA isolated from the tall revertants and dwarf plants with the *dw3-ref* allele. The PCR products were subsequently cloned and sequenced. The results obtained showed that a duplication of 882 bp had occurred in exon 5 of the *dw3* gene that led to the generation of the *dw3-ref* mutant allele (SEQ ID
20 NO: 1). Thus, the *dw3* dwarf phenotype in sorghum is likely due to an insertion-induced mutation that occurred within the *Dw3* allele to give rise to the *dw3-ref* allele. A partial sequence of the tall revertant allele, designated *Dw3-T* is disclosed in SEQ ID NO: 3. The duplication present in the *dw3-ref* mutant allele also seems to be responsible for the unstable nature of *dw3-ref*. By an undetermined mechanism, this duplication is removed
25 in tall revertants of *dw3-ref*.

Comparison of the partial amino acid sequence of the protein encoded by *Dw3-T* (SEQ ID NO: 4) revealed that, like BR2, this protein belongs to the family of multidrug-resistance-like P-glycoproteins. Whereas it shows more than 96% amino acid identity with the maize *pgp1* (the *Br2* gene), it exhibits 81% and 79% identity with P-glycoprotein genes of *Arabidopsis thaliana* and potato respectively.
30

Since the instability of the *dw3-ref* allele may result from some genetic recombination between two copies of the duplicated part of the gene, it might not always be precise. Some instances may occur where one or more extra base pairs may be left behind or deleted, leading in either case to a frame shift mutation. Such events are thus expected to generate new mutant alleles of *dw3* that are devoid of the duplication. And since the duplication seems to be responsible for the instability of *dw3-ref*, the new mutant alleles of *dw3* are expected to exhibit a stable dwarfing phenotype. Such stable dwarf alleles are highly desirable for breeding improved sorghum cultivars, as the instability of *dw3* has been a constant nemesis for breeders for enhancing the production of sorghum.

To identify a stable *dw3* allele, DNA was extracted from 200 dwarf sorghum plants and subjected to Southern blot analysis using a probe from the maize *Br2* gene. Two dwarf plants were identified that exhibited a restriction pattern that was different from the rest of the dwarf plants. Genomic DNA was isolated from one of these two dwarf plants and amplified using the oligonucleotide primers (SEQ ID NOS: 5-6) as described *supra*. The PCR product was cloned and sequenced. Comparison of the nucleotide sequence of the cloned PCR product (SEQ ID NO: 2) from this dwarf plant to the sequence of *dw3-ref* (SEQ ID NO: 1) revealed that the duplication present in *dw3-ref* was lost. Thus, this dwarf plant possesses a new *dw3* allele, designated as *dw3-1*. Comparison of the nucleotide sequence of the *dw3-1* allele with the *Dw3-T* allele demonstrated that the new *dw3-1* allele has undergone minor changes.

To separate the new *dw3-1* allele from the parental *dw3-ref* allele, the dwarf plant possessing the *dw3-1* was self pollinated and seeds from plant were collected and planted. From the progeny, plants that were homozygous for *dw3-1* were identified by Southern blot analysis, and the homozygous plants are being propagated to develop stable dwarfing germplasm for sorghum. In addition, eight separate Pioneer proprietary sorghum inbreds are also being genotyped for the presence of new mutant derivatives of *dw3-ref*. The inbreds that were utilized are AGK1G, MK7G, MQC100G, ZYL24, YYU28W, CAJ14W, FYL14W, and YGC87W. They were selected on the basis of their reversion frequency, which was rated high, moderate or low. These inbreds were planted outdoors in Johnston, Iowa during the summer of 1999. Two hundred plants from each line were

RFLP genotyped by digesting their DNA with PstI and hybridizing the resulting blots with a gene specific probe from the 3' end of the maize *br2* gene. Four stable homozygous dwarf plants were identified from YYU28W and ten such plants were identified from FYL14W. Seeds from these stable dwarf plants have been harvested.

- 5 The progeny of these stable dwarf plants can be used directly for the production of high-yielding sorghum hybrids with the desired stable dwarf phenotype.

EXAMPLE 2

Nucleotide Sequence of a *Dw3* Gene that Encodes a Functional Gene Product

10

- In order to clone the entire sequence from both the functional (*Dw3*) and the mutant (*dw3-ref*) alleles of the *dw3* locus, a tall revertant plant and a dwarf sibling were selected from the inbred AGK1G. In this inbred line, tall plants appear at a frequency of 0.1- 0.4 %. The genotype of the tall revertant plant was expected to be heterozygous at the *dw3* locus but otherwise identical to its dwarf sibling throughout the genome. To confirm that the tall revertant was heterozygous at the *dw3* locus, DNA samples isolated from this plant and a number of dwarf siblings were characterized by Southern analysis using three probes representing the 5', middle and 3' parts of the maize *br2* gene. As expected, polymorphism between dwarf siblings and the tall plant was localized only at the 3' end of *dw3*. This analysis allowed the identification of two EcoRI fragments from the tall revertant that when combined contained the entire *Dw3* allele. These were a 14kb EcoRI fragment that contained the 5' portion of the gene and an 8.1 kb fragment that contained the rest of the gene. A 9.0 kb fragment from the *dw3-ref* allele was determined to correspond to the 8.1 kb EcoRI fragment of the *Dw3* allele. Three size-selected libraries (containing the 14 kb/EcoRI and 8.1 kb/EcoRI fragments from the tall revertant and the 9.0 kb/EcoRI fragment from the dwarf sibling) were constructed in Lambda cloning vectors of Stratagene. The 14.0 kb /EcoRI fragment library was constructed in λ Dash II and was screened with a probe coming from the extreme 5' end of the maize *br2* gene. The other two libraries were prepared in λ ZapII and were screened with a probe from the 3' end of the maize *br2* gene. Positive clones were isolated and λ DNA was extracted for each of these clones. The *Dw3* and *dw3* genes were PCR amplified into
- 15
- 20
- 25
- 30

four overlapping 0.5 kb, 2.4 kb, 3.0 kb, and 1.3 kb fragments using gene specific primers and λ DNA as a template. These PCR fragments were cloned in TOPO vector (Invitrogen). From the dwarf *dw3* clone of 9.0 kb, a unique 888 bp *SacI* fragment containing a part of the duplicated region was subcloned into pBSK+ vector (Stratagene).

5 DNA from at least two colonies of each PCR clone was sequenced using M13 forward, M13 reverse, and gene specific primers (GSPs). The 888 bp *SacI* fragment from the *dw3*-ref clone was sequenced by using T3 and T7 vector-specific primers alone. Sequence information, both from the extreme 5' and 3' ends of *Dw3* and *dw3* genes, was gathered by sequencing directly the λ DNA of both the 14.0 kb and 8.1 kb clones, using
10 gene-specific primers. All of the sequence information was compiled and compared to reveal the cause of dwarfing in sorghum. A pairwise alignments between *Dw3* and *Br2* genes was done at the protein level by using Clustal W Program and at the nucleotide level by using BLAST Program of NCBL.

A polynucleotide of 6827 bp containing the full length *Dw3* gene was assembled
15 and is presented in SEQ ID NO: 7. Structurally, the *Dw3* gene has five exons and four introns. The length of five exons, from exon 1 through exon 5, is 616 bp, 537 bp, 326 bp, 230 bp, and 2400 bp, respectively. Intron 1 is 165 bp (nucleotides 639-803 of SEQ ID NO: 7); intron 2 is 110 bp (nucleotides 1441-1550); intron 3 is 846 bp (nucleotides 1877-2722); and intron 4 is 1471 bp (nucleotides 2953-4423) in length. The intron/exon
20 boundaries of *Dw3* are identical to that of the *br2* gene of maize. The predicted *Dw3*-cDNA is 4209 bp long from the start codon to the end of the termination codon (SEQ ID NO: 8) and is thus 28 bp longer than the analogous region of the *Br2*-cDNA. Similarly, the predicted protein encoded by *Dw3* is 1402 amino acids long (SEQ ID NO: 9), as compared to the 1394 amino acids predicted protein from *Br2* gene. Multiple alignment
25 results show that overall *Dw3* is 92% and 91.8% identical to the maize *Br2* gene at the nucleotide level and at the amino acid level, respectively.

PCR analysis of the polymorphic region between *dw3* and *br2* had earlier suggested that a duplication of a part of exon 5 resulted in the *dw3*-ref dwarfing allele of sorghum. To address if it was exclusively the reason for the mutant nature of the *dw3*-ref
30 allele, the sequence of the *Dw3* allele from the tall revertant was compared with that of the *dw3*-ref allele. As shown previously, the difference was detected only in exon 5

between these alleles. In the mutant allele (*dw3*-ref, SEQ ID NO: 1), a stretch of 882 bp in exon 5 (from nucleotides 5650-6531 of SEQ ID NO: 7) is duplicated at the 6532 nucleotide position in the same direction. This duplication converted the 1312 bp PstI restriction fragment (from nucleotides 5463 bp to 6775 of SEQ ID NO: 7) in the functional *Dw3* allele to the 2194 bp PstI fragment found in the *dw3*-ref allele, and thus was the cause for the polymorphism between these two alleles. Since no other changes were found between these alleles, the results clearly implicate this duplication as the sole cause for creating the *dw3* dwarfing allele of sorghum. The addition of 882 bp to the cDNA will no doubt have a serious ramification for the structure and activity of the DW3 protein. These findings also show how the *dw3*-ref allele spontaneously corrects itself, every now and then, by getting rid of the duplication. The mechanism, by which this correction occurs, remains unknown, as does the mechanism by which the duplication occurred in the first place.

Example 3

Transformation of Maize by Particle Bombardment and Regeneration of Transgenic Plants

Immature maize embryos from greenhouse donor plants are bombarded with a plasmid containing a P-glycoprotein nucleotide sequence of the invention operably linked to a promoter that drives expression in a plant and the selectable marker gene PAT (Wohlleben *et al.* (1988) *Gene* 70:25-37), which confers resistance to the herbicide Bialaphos. Alternatively, the selectable marker gene is provided on a separate plasmid. Transformation is performed as follows. Media recipes follow below.

Preparation of Target Tissue

The ears are husked and surface sterilized in 30% Clorox bleach plus 0.5% Micro detergent for 20 minutes, and rinsed two times with sterile water. The immature embryos are excised and placed embryo axis side down (scutellum side up), 25 embryos per plate, on 560Y medium for 4 hours and then aligned within the 2.5-cm target zone in preparation for bombardment.

Preparation of DNA

A plasmid vector comprising the P-glycoprotein nucleotide sequence of the invention operably linked to the plant promoter of interest is made. This plasmid DNA plus plasmid DNA containing a PAT selectable marker is precipitated onto 1.1 μm

5 (average diameter) tungsten pellets using a CaCl_2 precipitation procedure as follows:

100 μl prepared tungsten particles in water

10 μl (1 μg) DNA in Tris EDTA buffer (1 μg total DNA)

100 μl 2.5 M CaCl_2

10 μl 0.1 M spermidine

10 Each reagent is added sequentially to the tungsten particle suspension, while maintained on the multitube vortexer. The final mixture is sonicated briefly and allowed to incubate under constant vortexing for 10 minutes. After the precipitation period, the tubes are centrifuged briefly, liquid removed, washed with 500 ml 100% ethanol, and centrifuged for 30 seconds. Again the liquid is removed, and 105 μl 100% ethanol is
15 added to the final tungsten particle pellet. For particle gun bombardment, the tungsten/DNA particles are briefly sonicated and 10 μl spotted onto the center of each macrocarrier and allowed to dry about 2 minutes before bombardment.

Particle Gun Treatment

The sample plates are bombarded at level #4 in particle gun #HE34-1 or #HE34-
20 2. All samples receive a single shot at 650 PSI, with a total of ten aliquots taken from each tube of prepared particles/DNA.

Subsequent Treatment

Following bombardment, the embryos are kept on 560Y medium for 2 days, then transferred to 560R selection medium containing 3 mg/liter Bialaphos, and subcultured
25 every 2 weeks. After approximately 10 weeks of selection, selection-resistant callus clones are transferred to 288J medium to initiate plant regeneration. Following somatic embryo maturation (2-4 weeks), well-developed somatic embryos are transferred to medium for germination and transferred to the lighted culture room. Approximately 7-10 days later, developing plantlets are transferred to 272V hormone-free medium in tubes
30 for 7-10 days until plantlets are well established. Plants are then transferred to inserts in

flats (equivalent to 2.5" pot) containing potting soil and grown for 1 week in a growth chamber, subsequently grown an additional 1-2 weeks in the greenhouse, then transferred to classic 600 pots (1.6 gallon) and grown to maturity. Plants are monitored and scored for dwarf phenotype or other phenotype associated with expression of the P-glycoprotein nucleotides sequence of the invention.

Bombardment and Culture Media

Bombardment medium (560Y) comprises 4.0 g/l N6 basal salts (SIGMA C-1416), 1.0 ml/l Eriksson's Vitamin Mix (1000X SIGMA-1511), 0.5 mg/l thiamine HCl, 120.0 g/l sucrose, 1.0 mg/l 2,4-D, and 2.88 g/l L-proline (brought to volume with D-I H₂O following adjustment to pH 5.8 with KOH); 2.0 g/l Gelrite (added after bringing to volume with D-I H₂O); and 8.5 mg/l silver nitrate (added after sterilizing the medium and cooling to room temperature). Selection medium (560R) comprises 4.0 g/l N6 basal salts (SIGMA C-1416), 1.0 ml/l Eriksson's Vitamin Mix (1000X SIGMA-1511), 0.5 mg/l thiamine HCl, 30.0 g/l sucrose, and 2.0 mg/l 2,4-D (brought to volume with D-I H₂O following adjustment to pH 5.8 with KOH); 3.0 g/l Gelrite (added after bringing to volume with D-I H₂O); and 0.85 mg/l silver nitrate and 3.0 mg/l bialaphos (both added after sterilizing the medium and cooling to room temperature).

Plant regeneration medium (288J) comprises 4.3 g/l MS salts (GIBCO 11117-074), 5.0 ml/l MS vitamins stock solution (0.100 g nicotinic acid, 0.02 g/l thiamine HCL, 0.10 g/l pyridoxine HCL, and 0.40 g/l glycine brought to volume with polished D-I H₂O) (Murashige and Skoog (1962) *Physiol. Plant.* 15:473), 100 mg/l myo-inositol, 0.5 mg/l zeatin, 60 g/l sucrose, and 1.0 ml/l of 0.1 mM abscisic acid (brought to volume with polished D-I H₂O after adjusting to pH 5.6); 3.0 g/l Gelrite (added after bringing to volume with D-I H₂O); and 1.0 mg/l indoleacetic acid and 3.0 mg/l bialaphos (added after sterilizing the medium and cooling to 60°C). Hormone-free medium (272V) comprises 4.3 g/l MS salts (GIBCO 11117-074), 5.0 ml/l MS vitamins stock solution (0.100 g/l nicotinic acid, 0.02 g/l thiamine HCL, 0.10 g/l pyridoxine HCL, and 0.40 g/l glycine brought to volume with polished D-I H₂O), 0.1 g/l myo-inositol, and 40.0 g/l sucrose (brought to volume with polished D-I H₂O after adjusting pH to 5.6); and 6 g/l bacto-agar (added after bringing to volume with polished D-I H₂O), sterilized and cooled to 60° C.

Example 4

Agrobacterium-Mediated Transformation of Maize and Regeneration of Transgenic Plants

For *Agrobacterium*-mediated transformation of maize with a P-glycoprotein
5 nucleotide sequence of the invention, preferably the method of Zhao is employed (U.S.
Patent No. 5,981,840, and PCT patent publication WO98/32326; the contents of which are
hereby incorporated by reference). Briefly, immature embryos are isolated from maize and
the embryos contacted with a suspension of *Agrobacterium*, where the bacteria are
capable of transferring the P-glycoprotein nucleotide sequence of the invention to at least
10 one cell of at least one of the immature embryos (step 1: the infection step). In this step
the immature embryos are preferably immersed in an *Agrobacterium* suspension for the
initiation of inoculation. The embryos are co-cultured for a time with the *Agrobacterium*
(step 2: the co-cultivation step). Preferably the immature embryos are cultured on solid
medium following the infection step. Following this co-cultivation period an optional
15 "resting" step is contemplated. In this resting step, the embryos are incubated in the
presence of at least one antibiotic known to inhibit the growth of *Agrobacterium* without
the addition of a selective agent for plant transformants (step 3: resting step). Preferably
the immature embryos are cultured on solid medium with antibiotic, but without a
selecting agent, for elimination of *Agrobacterium* and for a resting phase for the infected
20 cells. Next, inoculated embryos are cultured on medium containing a selective agent and
growing transformed callus is recovered (step 4: the selection step). Preferably, the
immature embryos are cultured on solid medium with a selective agent resulting in the
selective growth of transformed cells. The callus is then regenerated into plants (step 5:
the regeneration step), and preferably calli grown on selective medium are cultured on
25 solid medium to regenerate the plants.

All publications and patent applications mentioned in the specification are
indicative of the level of those skilled in the art to which this invention pertains. All
publications and patent applications are herein incorporated by reference to the same
extent as if each individual publication or patent application was specifically and
30 individually indicated to be incorporated by reference.

Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be obvious that certain changes and modifications may be practiced within the scope of the appended claims.

5

RTA01/2085893v1

THAT WHICH IS CLAIMED:

1. An isolated nucleotide molecule comprising a nucleotide sequence selected from the group consisting of:

- 5 (a) a nucleotide sequence set forth in SEQ ID NO: 1;
- (b) a nucleotide sequence set forth in SEQ ID NO: 2;
- (c) a nucleotide sequence set forth in SEQ ID NO: 3;
- (d) a nucleotide sequence set forth in SEQ ID NO: 7;
- (e) a nucleotide sequence set forth in SEQ ID NO: 8;
- 10 (f) a nucleotide sequence consisting of at least 19 contiguous nucleotides of the nucleotide sequence set forth in any one of (a)-(e);
- (h) a nucleotide sequence encoding the amino acid sequence set forth in SEQ ID NO: 9;
- (i) a nucleotide sequence encoding at least 70 contiguous amino acids
- 15 of the amino acid sequence set forth in SEQ ID NO: 9;
- (j) a nucleotide sequence comprising at least 80% identity to the sequence set forth in SEQ ID NO: 7;
- (k) a nucleotide sequence comprising at least 80% identity to the sequence set forth in SEQ ID NO: 8;
- 20 (l) a nucleotide sequence that is complementary to the nucleotide sequence of any one of (a)-(k); and
- (m) a nucleotide sequence that hybridizes under stringent conditions to the nucleotide sequence any one of (a)-(e), or to a complementary sequence thereof.

25 2. An expression cassette comprising the nucleotide sequence of claim 1, said nucleotide sequence operably linked to a promoter that drives expression in a plant cell.

30 3. The expression cassette of claim 2, wherein said promoter is selected from the group consisting of tissue-preferred, constitutive, chemically regulatable, and pathogen-preferred promoters.

4. A transformed plant having stably incorporated into its genome a nucleotide sequence operably linked to a promoter that drives expression in a plant cell, wherein said nucleotide sequence is selected from the group consisting of:

- 5 (a) a nucleotide sequence set forth in SEQ ID NO: 1;
- (b) a nucleotide sequence set forth in SEQ ID NO: 2;
- (c) a nucleotide sequence set forth in SEQ ID NO: 3;
- (d) a nucleotide sequence set forth in SEQ ID NO: 7;
- (e) a nucleotide sequence set forth in SEQ ID NO: 8;
- 10 (f) a nucleotide sequence consisting of at least 19 contiguous nucleotides of the nucleotide sequence set forth in any one of (a)-(e);
- (h) a nucleotide sequence encoding the amino acid sequence set forth in SEQ ID NO: 9;
- (i) a nucleotide sequence encoding at least 70 contiguous amino acids
- 15 of the amino acid sequence set forth in SEQ ID NO: 9;
- (j) a nucleotide sequence comprising at least 80% identity to the sequence set forth in SEQ ID NO: 7;
- (k) a nucleotide sequence comprising at least 80% identity to the sequence set forth in SEQ ID NO: 8;
- 20 (l) a nucleotide sequence that is complementary to the nucleotide sequence of any one of (a)-(k); and
- (m) a nucleotide sequence that hybridizes under stringent conditions to the nucleotide sequence any one of (a)-(e), or to a complementary sequence thereof.

25 5. The plant of claim 4, wherein said promoter is selected from the group consisting of tissue-preferred, constitutive, chemically regulatable, and pathogen-preferred promoters.

30 6. The plant of claim 4, wherein said nucleotide sequence is operably linked to said promoter for the production of antisense transcripts.

7. The plant of claim 4, wherein said plant is a monocot.
8. The plant of claim 7, wherein said monocot is selected from the group consisting of maize, wheat, rice, Basmati rice, sorghum, rye, millet and barley.
- 5
9. The plant of claim 4, wherein said plant is a dicot.
10. The plant of claim 9, wherein said dicot is selected from the group consisting of soybeans, sunflowers, safflowers, alfalfa, Brassica sp., cotton, peanuts and
- 10 fruit trees.
11. Transformed seed of the plant of claim 4.
12. Transformed seed of the plant of claim 5.
- 15
13. Transformed seed of the plant of claim 6.
14. Transformed seed of the plant of claim 7.
- 20
15. Transformed seed of the plant of claim 8.
16. Transformed seed of the plant of claim 9.
17. Transformed seed of the plant of claim 10.
- 25
18. A method for modifying the growth of an organism, said method comprising transforming an organism with a nucleotide sequence encoding a P-glycoprotein wherein said P-glycoprotein functions to control growth of an organism, said nucleotide sequence operably linked to a promoter capable of driving the expression
- 30 of said sequence in said organism.

19. The method of claim 18, wherein said organism is a plant and said nucleotide sequence is selected from the group consisting of:

- (a) a nucleotide sequence set forth in SEQ ID NO: 1;
- (b) a nucleotide sequence set forth in SEQ ID NO: 2;
- 5 (c) a nucleotide sequence set forth in SEQ ID NO: 3;
- (d) a nucleotide sequence set forth in SEQ ID NO: 7;
- (e) a nucleotide sequence set forth in SEQ ID NO: 8;
- (f) a nucleotide sequence consisting of at least 19 contiguous nucleotides of the nucleotide sequence set forth in any one of (a)-(e);
- 10 (h) a nucleotide sequence encoding the amino acid sequence set forth in SEQ ID NO: 9;
- (i) a nucleotide sequence encoding at least 70 contiguous amino acids of the amino acid sequence set forth in SEQ ID NO: 9;
- (j) a nucleotide sequence comprising at least 80% identity to the
15 sequence set forth in SEQ ID NO: 7;
- (k) a nucleotide sequence comprising at least 80% identity to the sequence set forth in SEQ ID NO: 8;
- (l) a nucleotide sequence that is complementary to the nucleotide sequence of any one of (a)-(k); and
- 20 (m) a nucleotide sequence that hybridizes under stringent conditions to the nucleotide sequence any one of (a)-(e), or to a complementary sequence thereof.

20. The method of claim 18, wherein said nucleotide sequence is operably linked to said promoter for the production of antisense transcripts.

25

21. The method of claim 18, wherein the height of said plant is reduced.

22. The method of claim 18, wherein said plant is a monocot.

30

23. The method of claim 18, wherein said monocot is selected from the group consisting of maize, wheat, rice, Basmati rice, sorghum, rye, millet and barley.

24. A transformed plant cell having stably incorporated into its genome a nucleotide sequence operably linked to a promoter that drives expression in a plant cell, wherein said nucleotide sequence is selected from the group consisting of:

- 5 (a) a nucleotide sequence set forth in SEQ ID NO: 1;
- (b) a nucleotide sequence set forth in SEQ ID NO: 2;
- (c) a nucleotide sequence set forth in SEQ ID NO: 3;
- (d) a nucleotide sequence set forth in SEQ ID NO: 7;
- (e) a nucleotide sequence set forth in SEQ ID NO: 8;
- 10 (f) a nucleotide sequence consisting of at least 19 contiguous nucleotides of the nucleotide sequence set forth in any one of (a)-(e);
- (h) a nucleotide sequence encoding the amino acid sequence set forth in SEQ ID NO: 9;
- (i) a nucleotide sequence encoding at least 70 contiguous amino acids
- 15 of the amino acid sequence set forth in SEQ ID NO: 9;
- (j) a nucleotide sequence comprising at least 80% identity to the sequence set forth in SEQ ID NO: 7;
- (k) a nucleotide sequence comprising at least 80% identity to the sequence set forth in SEQ ID NO: 8;
- 20 (l) a nucleotide sequence that is complementary to the nucleotide sequence of any one of (a)-(k); and
- (m) a nucleotide sequence that hybridizes under stringent conditions to the nucleotide sequence any one of (a)-(e), or to a complementary sequence thereof.

25. A method for identifying a plant having a stable mutant phenotype comprising identifying a plant that lacks in its genome at least a portion of an insertion in a gene, wherein: said plant is a descendent of an unstable mutant plant which comprises in its genome an insertion in at least one copy of said gene; and said insertion in said gene is capable of causing said mutant phenotype.

30

26. The method of claim 25, wherein said insertion is a duplication or a transposon.

27. The method of claim 25, wherein said phenotype is a dwarf phenotype.

28. The method of claim 25, wherein said gene encodes a P-glycoprotein.

29. The method of claim 28, wherein said plant is sorghum and said gene is *Dw3*.

30. A plant having a stable mutant phenotype according to claim 25.

31. Seed of the plant of claim 30.

32. An isolated protein comprising a member selected from the group consisting of:

(a) a polypeptide comprising the amino acid sequence set forth in SEQ ID NO: 9;

(b) a polypeptide comprising at least 70 contiguous amino acids of the amino acid sequence set forth in SEQ ID NO: 9;

(c) a polypeptide encoded by the nucleotide sequence set forth in SEQ ID NO: 8; and

(d) a polypeptide encoded by an amino acid sequence comprising at least 80% identity to the amino sequence of (a).

SORGHUM DWARFING GENES AND METHODS OF USE

ABSTRACT

The invention relates to the genetic manipulation of organisms, particularly to the expression of P-glycoprotein genes in transformed plants and other organisms.

Nucleotide sequences for the P-glycoprotein genes, particularly the *Dw3* gene of
5 sorghum, and methods for their use are provided. The sequences find use in modifying the growth of organisms, particularly plants. Additionally, the invention provides methods for producing stable dwarf crop plants, particularly stable dwarf sorghum plants.

SEQUENCE LISTING

<110> Johal, Gurmukh S
Multani, Dilbag S

<120> SORGHUM DWARFING GENES AND METHODS OF USE

<130> 5718-100 (035718/205458)

<140>

<141>

<150> 60/165,176

<151> 1999-11-12

<160> 9

<170> PatentIn Ver. 2.1

<210> 1

<211> 2139

<212> DNA

<213> Sorghum bicolor

<400> 1

```
ctectcgccg tgttcccgt cgtcgtgggc gccaccgtgc tgcagaagat gttcatgaag 60
ggcttctcgg gggacctgga ggccgcgcac gccagggcca cgcagatcgc gggcgaggcc 120
gtggccaacc tgcgcaccgt ggccgcgttc aacgcggagc gcaagatcac ggggctgttc 180
gaggccaacc tgcgcggccc gctccggcgc tgcttctgga aggggcagat cgccggcagc 240
ggctacggcg tggcgcagtt cctgctgtac gogtccctac cgtcggggct gtggtacgcg 300
gcgtggctgg tgaagcacgg cgtgtccgac ttctcgcgca ccatccgcgt gttcatggtg 360
ctgatggtgt ccgccaacgg cgccgccgag acgctgacgc tggcgccgga ctttgtcaag 420
ggcgggcgcg cgatgcggtc cgtgttcgag accatcgacc ggaaaacgga ggtggagccc 480
gacgacgtgg acgcggcgcc ggtgccggag cggcccaagg gcgaggtgga gctgaagcac 540
gtggaacttct cgtacccgtc gcggccggac atccaggtgt tccgcgacct gagcctccgg 600
gcgcgcgcgg ggaagacgct ggcgctggtg ggtccgagcg ggtgcggcaa gagctcggtg 660
ctggcgctgg cgcagcggtt ctacgagccc acgtccgggc gcgtgctcct ggacggcaag 720
gacgtgcgca agtacaacct gcgggcgctg cggcgcgctg tggcggtggt gccgcaggag 780
ccgttccctgt tcgcggcgag catccacgac aacatcgcgt acgggcgcga gggcgcgacg 840
gaggcgaggg tgggtggaggc ggcgactcaa gcgaacgcgc accggttcat ctcggcgctg 900
ccggaggggct acgggacgca agtgggcgag cgcgggggtgc agctgtcggg cgggcagcgg 960
cagcggatcg cgatcgcgcg cgcgctggtg aagcaagcgg ccatcatgcc gctggacgag 1020
gcgaccagcg cgctggaccc gagtcggagc ggtggctctt cgaggccaac cttcgcggcc 1080
cgctccggcg cttgttctgg aaggggcaga tcgcgggaac ggtacgggcg tggcgcagtt 1140
cttgctgacg cgtcctacgc gcttgggggt ttggtacccc gcgtggctag tgaagcacgg 1200
gtctccgact ttgcgcacc atccgggtgt tcatggtgct gatggtgtcc gccaacggcg 1260
ccgccgagac gctgacgctg gcgcgggact ttgtcaaggg cgggcgcgcg atgcggtcgg 1320
tgttcgagac catcgaccgg aaaacggagg tggagcccga cgacgtggac gcggcgccgg 1380
```


tgccggagcg	gcccaggggc	gaggtggagc	tgaagcacgt	ggacttctcg	taccgcgtcg	1440
ggccggacat	ccaggtgttc	cgcgacctga	gcctccgggc	gcgcgccggg	aagacgctgg	1500
cgctggtggg	tccgagcggg	tgcggcaaga	gctcgggtgt	ggcgtggtg	cagcggttct	1560
acgagcccac	gtccggggcg	gtgctcctgg	acggcaagga	cgtgcgcaag	tacaacctgc	1620
gggcgtgctg	gcgcgtggtg	gcggtggtgc	cgcaggagcc	gttcctgttc	gcggcgagca	1680
tccacgacaa	catcgcgtag	gggcgcgagg	gcgcgacgga	ggcggagggtg	gtggaggcgg	1740
cgacgcaggc	gaacgcgcac	cggttcatct	cggcgtgtcc	ggagggctac	gggacgcagg	1800
tgggcgagcg	cgggggtgcag	ctgtcggggc	ggcagcggca	gcggatcgcg	atcgcgcgcg	1860
ctggtaaagca	gcggccatcg	tgtgtctgga	cgaggcgacc	agcgcgctgg	acgccgagtc	1920
ggagcgggtgc	gtgcaggagg	cgctggagcg	cgcgggggtcc	gggcgcacca	ccatcgtggt	1980
ggcgaccggg	ctggccacgg	tgcgcggcg	gcacaccatc	gcggtcatcg	acgacggcaa	2040
ggtggcggag	caggggtcgc	actcgcacct	gctcaagcac	catcccgacg	ggtgctacgc	2100
gcggatgctg	cagctgcagc	ggctgacggg	cgcggcggc			2139

<210> 2

<211> 1267

<212> DNA

<213> Sorghum bicolor

<400> 2

ctcctcgccg	tgttcccgt	cgctcgtgggc	gccaccgtgc	tgcagaagat	gttcatgaag	60
ggcttctcgg	gggacctgga	ggccgcgcac	gccagggcca	cgcagatcgc	gggcgaggcc	120
gtggccaacc	tgcgcaccgt	ggccgcgttc	aacgcggagc	gcaagatcac	ggggctgttc	180
gaggccaacc	tgcgcggccc	gctccggcgc	tgttcttgga	aggggcagat	cgccggcagc	240
ggctacggcg	tggcgcagtt	cctgctgtac	gcgtcctacg	cgtggggct	gtggtacgcg	300
gcgtggctgg	tgaagcacgg	cgtgtccgac	ttctcgcgca	ccatccgcgt	gttcatggtg	360
ctgatggtgt	ccgcgaacgg	gcgccgcgcg	agacgtgac	gctggcgccg	gacttcatca	420
agggcggggc	cgcgatgcgg	tcggtgttcg	agacgatcga	cgcgaagacg	gaggtggagc	480
ccgacgacgt	ggacgcggcg	ccggtgccgg	agcggccgag	gggcgagggtg	gagctgaagc	540
acgtggactt	ctcgtacccg	tcgcggccgg	acatccaggt	gttccgcgac	ctgagcctcc	600
gtgcgcgcgc	cgggaagacg	ctggcgctgg	tggggccgag	cgggtgcggc	aagagctcgg	660
tgctggctct	ggtgcagcgg	ttctacaagc	ccacgtccgg	gcgcgtgttc	ttgacggcaa	720
agacgtgcgc	aaaaacaacc	ttcgggcgtt	ccggcgcatt	gttgcggtgg	taccaagaa	780
cccgtttcct	gttcgcggcg	aagaatccac	gagaacatcg	cgcacggggc	agagggcgct	840
acggaggcgg	aggtggtgga	ggcggcgggc	caggcgaacg	cgcaccgggtt	catcgcgggc	900
ctgccggaag	gggtactgga	cgcagaaggg	cgcgcgcggg	gtgcacctgt	cggggggcag	960
cggcagcgga	tcgcgatcgc	gcgcgcgctg	gtgaagcagc	ggccatcgtg	ctgctggacg	1020
agggcaccag	cgcgctggac	gccgagtcgg	agcgggtgcgt	gcaggaggcg	ctggagcgcg	1080
cggggtcggg	gcgcaccacc	atcgtggtgg	cgcaccggct	ggccacgggtg	cgcggcgcg	1140
acaccatcgc	ggtcatcgac	gacggcaagg	tggcggagca	ggggtcgcac	tcgcacctgc	1200
tcaagcacca	tcccgacggg	tgtacgcgc	ggatgctgca	gctgcagcgg	ctgacggggc	1260
cggcggc						1267

<210> 3

<211> 1261

<212> DNA

<213> Sorghum bicolor

<220>

<221> CDS

<222> (1)..(1245)

<400> 3

ctc ctc gcc gtg ttc ccg ctc gtc gtg ggc gcc acc gtg ctg cag aag	48
Leu Leu Ala Val Phe Pro Leu Val Val Gly Ala Thr Val Leu Gln Lys	
1 5 10 15	
atg ttc atg aag ggc ttc tcg ggg gac ctg gag gcc gcg cac gcc agg	96
Met Phe Met Lys Gly Phe Ser Gly Asp Leu Glu Ala Ala His Ala Arg	
20 25 30	
gcc acg cag atc gcg ggc gag gcc gtg gcc aac ctg cgc acc gtg gcc	144
Ala Thr Gln Ile Ala Gly Glu Ala Val Ala Asn Leu Arg Thr Val Ala	
35 40 45	
gcg ttc aac gcg gag cgc aag atc acg ggg ctg ttc gag gcc aac ctg	192
Ala Phe Asn Ala Glu Arg Lys Ile Thr Gly Leu Phe Glu Ala Asn Leu	
50 55 60	
cgc ggc ccg ctc cgg cgc tgc ttc tgg aag ggg cag atc gcc ggc agc	240
Arg Gly Pro Leu Arg Arg Cys Phe Trp Lys Gly Gln Ile Ala Gly Ser	
65 70 75 80	
ggc tac gcc gtg gcg cag ttc ctg ctg tac gcg tcc tac gcg ctg ggg	288
Gly Tyr Gly Val Ala Gln Phe Leu Leu Tyr Ala Ser Tyr Ala Leu Gly	
85 90 95	
ctg tgg tac gcg gcg tgg ctg gtg aag cac ggc gtg tcc gac ttc tcg	336
Leu Trp Tyr Ala Ala Trp Leu Val Lys His Gly Val Ser Asp Phe Ser	
100 105 110	
cgc acc atc cgc gtg ttc atg gtg ctg atg gtg tcc gcc aac ggc gcc	384
Arg Thr Ile Arg Val Phe Met Val Leu Met Val Ser Ala Asn Gly Ala	
115 120 125	
gcc gag acg ctg acg ctg gcg ccg gac ttt gtc aag ggc ggg cgc gcg	432
Ala Glu Thr Leu Thr Leu Ala Pro Asp Phe Val Lys Gly Gly Arg Ala	
130 135 140	
atg cgg tcc gtg ttc gag acc atc gac cgg aaa acg gag gtg gag ccc	480
Met Arg Ser Val Phe Glu Thr Ile Asp Arg Lys Thr Glu Val Glu Pro	
145 150 155 160	
gac gac gtg gac gcg gcg ccg gtg ccg gag cgg ccc aag ggc gag gtg	528

Asp Asp Val Asp Ala Ala Pro Val Pro Glu Arg Pro Lys Gly Glu Val	
165 170 175	
gag ctg aag cac gtg gac ttc tcg tac ccg tcg cgg ccg gac atc cag	576
Glu Leu Lys His Val Asp Phe Ser Tyr Pro Ser Arg Pro Asp Ile Gln	
180 185 190	
gtg ttc cgc gac ctg agc ctc cgg gcg cgc gcc ggg aag acg ctg gcg	624
Val Phe Arg Asp Leu Ser Leu Arg Ala Arg Ala Gly Lys Thr Leu Ala	
195 200 205	
ctg gtg ggt ccg agc ggg tgc ggc aag agc tcg gtg ctg gcg ctg gtg	672
Leu Val Gly Pro Ser Gly Cys Gly Lys Ser Ser Val Leu Ala Leu Val	
210 215 220	
cag cgg ttc tac gag ccc acg tcc ggg cgc gtg ctc ctg gac ggc aag	720
Gln Arg Phe Tyr Glu Pro Thr Ser Gly Arg Val Leu Leu Asp Gly Lys	
225 230 235 240	
gac gtg cgc aag tac aac ctg cgg gcg ctg cgg cgc gtg gtg gcg gtg	768
Asp Val Arg Lys Tyr Asn Leu Arg Ala Leu Arg Arg Val Val Ala Val	
245 250 255	
gcg ccg cag gag ccg ttc ctg ttc gcg gcg agc atc cac gac aac atc	816
Ala Pro Gln Glu Pro Phe Leu Phe Ala Ala Ser Ile His Asp Asn Ile	
260 265 270	
gcg tac ggg cgc gag ggc gcg acg gag gcg gag gtg gtg gag gcg gcg	864
Ala Tyr Gly Arg Glu Gly Ala Thr Glu Ala Glu Val Val Glu Ala Ala	
275 280 285	
acg cag gcg aac gcg cac cgg ttc atc gcg gcg ctg ccg gag ggc tac	912
Thr Gln Ala Asn Ala His Arg Phe Ile Ala Ala Leu Pro Glu Gly Tyr	
290 295 300	
ggg acg cag gtg ggc gag cgc ggg gtg cag ctg tcg ggc ggg cag ccg	960
Gly Thr Gln Val Gly Glu Arg Gly Val Gln Leu Ser Gly Gly Gln Arg	
305 310 315 320	
cag cgg atc gcg atc gcg cgc gct ggt aag cag cgg cca tcg tgc tgc	1008
Gln Arg Ile Ala Ile Ala Arg Ala Gly Lys Gln Arg Pro Ser Cys Cys	
325 330 335	
tgg acg agg cga cca gcg cgc tgg acg ccg agt cgg agc ggt gcg tgc	1056
Trp Thr Arg Arg Pro Ala Arg Trp Thr Pro Ser Arg Ser Gly Ala Cys	
340 345 350	
agg agg cgc tgg agc gcg cgg ggt ccg ggc gca cca cca tcg tgg tgg	1104

Arg Arg Arg Trp Ser Ala Arg Gly Pro Gly Ala Pro Pro Ser Trp Trp
355 360 365

cgc acc ggc tgg cca cgg tgc gcg gcg cgc aca cca tcg cgg tca tcg 1152
Arg Thr Gly Trp Pro Arg Cys Ala Ala Arg Thr Pro Ser Arg Ser Ser
370 375 380

acg acg gca agg tgg cgg agc agg ggt cgc act cgc acc tgc tca agc 1200
Thr Thr Ala Arg Trp Arg Ser Arg Gly Arg Thr Arg Thr Cys Ser Ser
385 390 395 400

acc atc ccg acg ggt gct acg cgc gga tgc tgc agc tgc agc ggc 1245
Thr Ile Pro Thr Gly Ala Thr Arg Gly Cys Cys Ser Cys Ser Gly
405 410 415

tgacgggccc ggcggc 1261

<210> 4
<211> 415
<212> PRT
<213> Sorghum bicolor

<400> 4
Leu Leu Ala Val Phe Pro Leu Val Val Gly Ala Thr Val Leu Gln Lys
1 5 10 15

Met Phe Met Lys Gly Phe Ser Gly Asp Leu Glu Ala Ala His Ala Arg
20 25 30

Ala Thr Gln Ile Ala Gly Glu Ala Val Ala Asn Leu Arg Thr Val Ala
35 40 45

Ala Phe Asn Ala Glu Arg Lys Ile Thr Gly Leu Phe Glu Ala Asn Leu
50 55 60

Arg Gly Pro Leu Arg Arg Cys Phe Trp Lys Gly Gln Ile Ala Gly Ser
65 70 75 80

Gly Tyr Gly Val Ala Gln Phe Leu Leu Tyr Ala Ser Tyr Ala Leu Gly
85 90 95

Leu Trp Tyr Ala Ala Trp Leu Val Lys His Gly Val Ser Asp Phe Ser
100 105 110

Arg Thr Ile Arg Val Phe Met Val Leu Met Val Ser Ala Asn Gly Ala
115 120 125

Ala	Glu	Thr	Leu	Thr	Leu	Ala	Pro	Asp	Phe	Val	Lys	Gly	Gly	Arg	Ala	130	135	140
Met	Arg	Ser	Val	Phe	Glu	Thr	Ile	Asp	Arg	Lys	Thr	Glu	Val	Glu	Pro	145	150	155
Asp	Asp	Val	Asp	Ala	Ala	Pro	Val	Pro	Glu	Arg	Pro	Lys	Gly	Glu	Val	165	170	175
Glu	Leu	Lys	His	Val	Asp	Phe	Ser	Tyr	Pro	Ser	Arg	Pro	Asp	Ile	Gln	180	185	190
Val	Phe	Arg	Asp	Leu	Ser	Leu	Arg	Ala	Arg	Ala	Gly	Lys	Thr	Leu	Ala	195	200	205
Leu	Val	Gly	Pro	Ser	Gly	Cys	Gly	Lys	Ser	Ser	Val	Leu	Ala	Leu	Val	210	215	220
Gln	Arg	Phe	Tyr	Glu	Pro	Thr	Ser	Gly	Arg	Val	Leu	Leu	Asp	Gly	Lys	225	230	235
Asp	Val	Arg	Lys	Tyr	Asn	Leu	Arg	Ala	Leu	Arg	Arg	Val	Val	Ala	Val	245	250	255
Ala	Pro	Gln	Glu	Pro	Phe	Leu	Phe	Ala	Ala	Ser	Ile	His	Asp	Asn	Ile	260	265	270
Ala	Tyr	Gly	Arg	Glu	Gly	Ala	Thr	Glu	Ala	Glu	Val	Val	Glu	Ala	Ala	275	280	285
Thr	Gln	Ala	Asn	Ala	His	Arg	Phe	Ile	Ala	Ala	Leu	Pro	Glu	Gly	Tyr	290	295	300
Gly	Thr	Gln	Val	Gly	Glu	Arg	Gly	Val	Gln	Leu	Ser	Gly	Gly	Gln	Arg	305	310	315
Gln	Arg	Ile	Ala	Ile	Ala	Arg	Ala	Gly	Lys	Gln	Arg	Pro	Ser	Cys	Cys	325	330	335
Trp	Thr	Arg	Arg	Pro	Ala	Arg	Trp	Thr	Pro	Ser	Arg	Ser	Gly	Ala	Cys	340	345	350
Arg	Arg	Arg	Trp	Ser	Ala	Arg	Gly	Pro	Gly	Ala	Pro	Pro	Ser	Trp	Trp	355	360	365
Arg	Thr	Gly	Trp	Pro	Arg	Cys	Ala	Ala	Arg	Thr	Pro	Ser	Arg	Ser	Ser	370	375	380

Thr Thr Ala Arg Trp Arg Ser Arg Gly Arg Thr Arg Thr Cys Ser Ser
 385 390 395 400

Thr Ile Pro Thr Gly Ala Thr Arg Gly Cys Cys Ser Cys Ser Gly
 405 410 415

<210> 5

<211> 26

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:oligonucleotide
 primer designed from sequence of Zea mays Br2 gene

<400> 5

ctcctcgccg tggtcccgt cgtcgt

26

<210> 6

<211> 17

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:oligonucleotide
 primer designed from sequence of Zea mays Br2 gene

<400> 6

gccgccgcgc ccgtcag

17

<210> 7

<211> 6827

<212> DNA

<213> Sorghum bicolor

<400> 7

gtccctcccc ggccccgat cgatgtctac caacgaccgc gacgagatca gggcgcgcggt 60
 cgtcgtcctc ggcgcccctc atgccgacga cgacgcgggc gacgagtggg ccgccccga 120
 gctcgaggcc ttccacctcc cctctcccgc ccaccagcct cctggcttcc acctagccgc 180
 tgggcaccaa ccggaagctg cagcagagca acccaccacg ctccctgctg ccgcccgcac 240
 cagcgacaca tccactgctg ctggtgctgc tctctcttct ccttcgcccgc ctccgcccgc 300
 ggctcctttg gagatggacc agccgcccaa tgccaagccg gcctcctcct ccgcccgcgc 360
 cgccggcgcc aatgacaaca agaagcccac ccgcccgcgc gcgctgcgcg acctcttccg 420
 cttcgccgac ggctcgact gcgcgctcat gctcgtcggc acgctcggcg cgctcgtcca 480

cggtctgctcg ctccccgtct tectccgctt cttecgccgac ctcgtcgact ccttcggctc 540
 ccacgccaac gacccggaaca ccatgggtccg cctcgtcgtc aagtacgcct tctacttctc 600
 cgctcgtcga gccgcaatct gggcgctctc atgggcaggt aaccaacgtt attcctcctc 660
 ctctccccct cctcctctcc ggcactgctg ctccgctcgc gaattgtctg tcgatttgga 720
 ttggatggcg aatcacatca gtcgctcaat cttecatggcc catggctagc aatgagatcg 780
 accttcgaat cctcgtcttg cagagatctc ctgctggatg tggaccggcg agcggcagtc 840
 gacgcggatg cggatccggg acctggacgc ggcgctgcgg caggacgtgt ccttcttcga 900
 caccgacgtg cgcacctcgg acgtcatcta cgccatcaac gcggacgccg tgggtgggtgc 960
 aggacgccat cagcgagaag ctgggcaacc tcatccacta catggccacc ttcgtggcgg 1020
 gcttcgtcgt gggcttcacc gccgcctggc agctggcgct cgtcacgctc gccgtcgtgc 1080
 cgctcatcgc cgtcatcggg gggctcagcg ccgcgcgcgt cgccaagctc tctccagga 1140
 gccaggacgc gctgtcgggc gccagcggca tcgcggagca ggcgctcgcg cagatacga 1200
 tcgtgcaggg cttcgtcggc gaggagcgcg aaatgcgggc gtactcggcg gcgttgggcg 1260
 tcgcgcagaa gatcggctac cgcagcggct tcgccaaggg gctcggcctc ggcggcacct 1320
 acttcaccgt cttctgctgc tacggcctcc tgctctggta cggcggacac ctcgtccgcg 1380
 gaaccacacc aacggagggc tcgccatcgc caccatgttc tccgtcatga tcggcgggct 1440
 gtaagatgat cagtttctcc cgggctctcc tgttcttccg tcatgacaca gcatgtacta 1500
 cgtacgctta ctgggtctgtg tctgtgtgtg tgtggatcgc ctgcgtccag ggccctcggg 1560
 cagtcggcgc cgagcatggc cgcgttcgcc aaggcgcgcg tggcggccgc caagatcttc 1620
 cgcacatcgc accacaggcc gggcatctcc tcgcgggacg gcgaggacgg cggcggcggtg 1680
 gagctggagt cgggtgacggg gcgggtggag atgaggggcg tggacttcgc gtaccgcgtc 1740
 cggccggacg tccccatcct gcgcggcttc tcgctcagcg tgcccgcggg caagaccatc 1800
 gcgctgggtg gcagctccgg ctccgggaag agcacgggtg tgcgtctcct cgagagggtc 1860
 tacgacccca gcgcaggtat acatagtagc ctaccaatc tagctttagc gcattgatta 1920
 attagtgttg gagttcactt gcttgccaat tgccattgcc atcacacatc agcagctacc 1980
 atacattgcc aactgccatt gctgctgcct tgctgggtgg ttagtagggg aagaagcttc 2040
 cactgtagca ggagtacatt gcaaacagga agtgaatttt gcacgtggga aatgaagaag 2100
 tgaatgcttg gagcagagct ggccggcctc atgggctgct tacctactat ctagtcaacc 2160
 aagcatccct gtttcttctc tgtttatggg caaggcattc acaccagctt agaaacttag 2220
 aaagaagcta agccctttgt tttaattttc ctaaaaaaaa atttttgggtt aaaatttttt 2280
 taaagtttcc cttgtccaca atcccaaatt cttttaagaa gtattaatat atgacgaaaa 2340
 ttaaaaccta attgcaccag ttttggtcga aattgaccga aaacgatctt ttgagcctaa 2400
 ttagtcccat gattggacaa tatttgtcaa atacaaacga aagtgtctgc gtatcgattt 2460
 tgcaaagggt ttcggaacta aacaaagcct ggtgcctgca acgcgagaca aagaaaacta 2520
 tttgcctggc aagatgccac tattgcacat gcatgccact ctttgagcct tgaccgactg 2580
 actgactact cagagtagga gtgggttcaat tgtattgaca ttagtagga gtactcgtat 2640
 gctatagtag tctgttagct ttttcaaaca aaaaaaaaag agaaagaaag aatgaagtc 2700
 tgaaatttgt tgggttttggc agggcaaata ttgctggacg ggcattgatc caagtcgctg 2760
 aagctccggg ggctccggca gcagattggg ctggtgagcc aggagccgac gctgttcgcg 2820
 acgagcatca aggagaacct gctgctgggg cgggacagtc agagtgcgac gcaggccgag 2880
 atggaggagg ccgccagggt ggccaacgcg cactccttca tcgtcaagct ccccgacggc 2940
 tacgacacgc aggtccgtcc cgtatagcta gctcactagc tgactgcca cttctctcgc 3000
 ttgctcccca ccgttgctgc ctggtgctct ccaatccact tgctgggtgc tggaccacac 3060
 gtgcctgctt gcctagctgc tccacatctg ctttccctgt ccaaccttat gcaactcact 3120
 ctaatactat atcaaataca tttctagagt ttaaagctta tcttagaata aatgcattct 3180
 tagctacgag acaacctaac ttcagttggt gttgttggtt tttttacttt ctctcttctc 3240
 acaaatacta tgattacgtc tttacagcga tcttttttat tccaaaccta aaaatgcatg 3300
 cactcactct aaaagcgcaa agggagcatc ttttttccc ccatcatctg cacgcagcct 3360

```

tttcttttcc tcatgtcacg agggactgaa ggtgtgtatg cagcgtcaag tcatccatcc 3420
gttccacttg gattaggacg ggggggccaa tttttaggcc cttgtttgcc attcgcatth 3480
tggtgtgttg ggtttgccaa caagaaactt gccaggttgc ttttgttatc acgcacagga 3540
caggagaggt ctttttctcg acacaagctc tacagcctct actaaactag cacttgctga 3600
tgagagcaga ggatgaatgg acgatgaaca tctagagtga gagagaaaaa aatgttaata 3660
ataataaaaa gtagtagcag gattaagaat caacctgggg tacgtaggaa gaggtacaat 3720
ccctaggaat ctagagtatg agaagtatgg gaggagttag gggagtggaa cggaacaaat 3780
tccgagttgg tattttgccg ggaatgtcaa gttgattttt gatcctagtg caagcaagaa 3840
ttatcaatca ctcagactca gcctgtctgt gtctgtccac cccagctctt gctactctac 3900
ttactactgt gctactagtg gtagggtagg tatcttacat aaactgttat tataaactgt 3960
catctgagaa agagagccag tcaaaccat gctgctgctt attttaatca ctgtcaaagt 4020
gcaggcaggc aggcagtctg gttagttaat aacatctggg aagggtttta tcaaaccaaa 4080
tcaaatacaga cgaaatctag aggccacatg ggatggggcc atatgtactg tactagcata 4140
actagcggct agatttttatt agaacacgga ctcacactcc cataactata actgacttga 4200
tcatgattcc ttgccaagca atgctcgcac gcccatgcat gcatcatccc tgggtcaaact 4260
caaacactct ccaccgtcag ggaataagac ttattatttt attaacaatt caatttttat 4320
ttattaatta cgtctggacg aggagtactg gtttatttga tgagagacat ggcagtccaa 4380
gtcaaactcg tttgtctgac catggcgggtg atggccgggtg cagggttgggg agcgcgccct 4440
gcagctctcc ggtgggcaga agcagcgcac cgccatcgcc cgcccatgac tcaagaacct 4500
cgccatcctg ctgctggacg aggccaccag cgcgctggac tccgagtctg agaagctcgt 4560
gcaggaggcg ctggaccgct tcatgatcgg gcgcaccacc ctggtgatcg cgcacaggat 4620
gtccaccatc cgcaaggcgg acgtggtggc cgtgctgcag ggcggccccg tctccgagat 4680
gggcgcgcac gacgagttga tggccaaggg cgagaacggc acttacgcca agttcatccg 4740
catgcaggag caggcgcacg aggcggcggt cgtcaacgcc cgccgcagca gcgccaggcc 4800
ctccagcgcc cgcaactcgg tcagctcgcc catcatgacg cgcaactcct cctacggccg 4860
ctccccatac tcccgcgcgc tctccgactt ctccacctcc gacttcaccc tctccatcca 4920
cgaccgcgac caccaccacc ggacgatggc cgacaagcag ctccgcttcc gcgcggcgcc 4980
cagctccttc ctccgcctcg ccaggatgaa ctccgcccag tgggcctacg cgctcgctcg 5040
ctccctgggc tccatggtct gcggctcctt cagcgccatc ttgcctaca tcctcagcgc 5100
cgtgctcagc gtctactacg cgccggaccc tcgctacatg aagcgcgaga tcgccaagta 5160
ctgctacctg ctcatcgcca tgtcctccgc ggcgctgctg ttcaacacgg tgcagcacgt 5220
gttctgggac acggtcggcg agaacctcac gaagcgtgtg cgcgagaaga tgttcgcccgc 5280
cgtgctccgc aacgagatcg cctggttcga cgccgacgag aacgccagcg cgcgcgctgc 5340
cgccaggctc gcgctcgacg cccagaacgt gcgctccgcc atcggggacc gtatctccgt 5400
catcgtccag aactcggcgc tcatgctcgt cgctcgacc gcgggcttcg tcctccagtg 5460
gcgcctcgcg ctctgtgctc tcgcccgtct cccgctcgtc gtggccgcca ccgtgctgca 5520
gaagatgttc atgaagggct tctcggggga cctggaggcc gcgcacgcca gggccacgca 5580
gatcgcgggc gaggccgtgg ccaacctgcg caccgtggcc gcgttcaacg cggagcgcaa 5640
gatcacgggg ctgttcgagg ccaacctgcg cgcccgcctc cgcgctgct tctggaaggg 5700
gcagatcgcc ggcagcggct acggcgtggc gcagttcctg ctgtacgcgt cctacgcgct 5760
ggggctgtgg tacgcggcgt ggctggtgaa gcacggcgtg tccgacttct cgcgccacct 5820
ccgcgtgttc atggtgctga tgggtgtccgc caacggcgcc gccgagacgc tgacgctggc 5880
gccggacttt gtcaagggcg ggcgcgcgat gcggtccgtg ttcgagacca tcgaccggaa 5940
aacggaggtg gagcccgacg acgtggacgc ggcgcgggtg ccggagcggc ccaaggcgca 6000
ggtggagctg aagcacgtgg acttctcgta cccgtcgcgg ccggacatcc aggtgttccg 6060
cgacctgagc ctccggggcg gcgcggggaa gacgctggcg ctggtgggtc cgagcgggtg 6120
cggcaagagc tcggtgctgg cgctggtgca gcggttctac gagccacgt ccgggcgcgt 6180
gctcctggac ggcaaggacg tgcgcaagta caacctgcgg gcgctgcggc gcgtggtggc 6240

```

```

ggtggcgccg caggagccgt tcctgttcgc ggcgagcatc cacgacaaca tcgcgtacgg 6300
gcgcgagggc gcgacggagg cggaggtggt ggaggcggcg acgcaggcga acgcgcaccg 6360
gttcatcgcg gcgctgccgg agggctacgg gacgcaggtg ggcgagcgcg ggggtgcagct 6420
gtcggggcggg cagcggcagc ggatcgcgat cgcgcgcgcg ctggtgaagc aggcggccat 6480
cgtgctgctg gacgaggcga ccagcgcgct ggacgccgag tcggagcggg gcgtgcagga 6540
ggcgctggag cgcgcggggg ccgggcgcac caccatcgtg gtggcgcacc ggctggccac 6600
ggtgcgcggc gcgcacacca tcgcggtcat cgacgacggc aagggtggcg agcagggggtc 6660
gcactcgcac ctgctcaagc accatcccga cgggtgctac gcgcggatgc tgcagctgca 6720
gcggctgacg ggcgggtgcc gcgccggggc cgccgccgtc gtcgtccaac ggggcccgcg 6780
cgtaggatgg atggatggat catggatgag tttggttct tgataaa 6827

```

<210> 8

<211> 4213

<212> DNA

<213> Sorghum bicolor

<220>

<221> CDS

<222> (1)..(4206)

<400> 8

```

atg tct acc aac gac ccg gac gag atc agg gcg cgc gtc gtc gtc ctc 48
Met Ser Thr Asn Asp Pro Asp Glu Ile Arg Ala Arg Val Val Val Leu
1 5 10 15

```

```

ggc gcc cct cat gcc gac gac gac gcc ggc gac gag tgg gcc cgc ccc 96
Gly Ala Pro His Ala Asp Asp Asp Ala Gly Asp Glu Trp Ala Arg Pro
20 25 30

```

```

gag ctc gag gcc ttc cac ctc ccc tct ccc gcc cac cag cct cct ggc 144
Glu Leu Glu Ala Phe His Leu Pro Ser Pro Ala His Gln Pro Pro Gly
35 40 45

```

```

ttc cac cta gcc gct ggg cac caa ccg gaa gct gca gca gag caa ccc 192
Phe His Leu Ala Ala Gly His Gln Pro Glu Ala Ala Ala Glu Gln Pro
50 55 60

```

```

acc acg ctc cct gct gcc cgc cgc acc agc gac aca tcc act gct gct 240
Thr Thr Leu Pro Ala Ala Arg Arg Thr Ser Asp Thr Ser Thr Ala Ala
65 70 75 80

```

```

ggt gct gct cct cct tct cct tcg ccg cct ccg ccg ccg gct cct ttg 288
Gly Ala Ala Pro Pro Ser Pro Ser Pro Pro Pro Pro Pro Ala Pro Leu
85 90 95

```

```

gag atg gac cag ccg ccc aat gcc aag ccg gcc tcc tcc tcc gcc gcc 336
Glu Met Asp Gln Pro Pro Asn Ala Lys Pro Ala Ser Ser Ser Ala Ala

```

100	105	110	
gcc gcc ggc gcc aat gac aac aag aag ccc acc ccg ccc gcc gcg ctg			384
Ala Ala Gly Ala Asn Asp Asn Lys Lys Pro Thr Pro Pro Ala Ala Leu			
115	120	125	
cgc gac ctc ttc cgc ttc gcc gac ggc ctc gac tgc gcg ctc atg ctc			432
Arg Asp Leu Phe Arg Phe Ala Asp Gly Leu Asp Cys Ala Leu Met Leu			
130	135	140	
gtc ggc acg ctc ggc gcg ctc gtc cac ggc tgc tcg ctc ccc gtc ttc			480
Val Gly Thr Leu Gly Ala Leu Val His Gly Cys Ser Leu Pro Val Phe			
145	150	155	160
ctc cgc ttc ttc gcc gac ctc gtc gac tcc ttc ggc tcc cac gcc aac			528
Leu Arg Phe Phe Ala Asp Leu Val Asp Ser Phe Gly Ser His Ala Asn			
165	170	175	
gac ccg gac acc atg gtc cgc ctc gtc gtc aag tac gcc ttc tac ttc			576
Asp Pro Asp Thr Met Val Arg Leu Val Val Lys Tyr Ala Phe Tyr Phe			
180	185	190	
ctc gtc gtc gga gcc gca atc tgg gcg tcc tca tgg gca gag atc tcc			624
Leu Val Val Gly Ala Ala Ile Trp Ala Ser Ser Trp Ala Glu Ile Ser			
195	200	205	
tgc tgg atg tgg acc ggc gag cgg cag tcg acg cgg atg cgg atc cgg			672
Cys Trp Met Trp Thr Gly Glu Arg Gln Ser Thr Arg Met Arg Ile Arg			
210	215	220	
tac ctg gac gcg gcg ctg cgg cag gac gtg tcc ttc ttc gac acc gac			720
Tyr Leu Asp Ala Ala Leu Arg Gln Asp Val Ser Phe Phe Asp Thr Asp			
225	230	235	240
gtg cgc acc tcg gac gtc atc tac gcc atc aac gcg gac gcc gtg gtg			768
Val Arg Thr Ser Asp Val Ile Tyr Ala Ile Asn Ala Asp Ala Val Val			
245	250	255	
ggt gca gga cgc cat cag cga gaa gct ggg caa cct cat cca cta cat			816
Gly Ala Gly Arg His Gln Arg Glu Ala Gly Gln Pro His Pro Leu His			
260	265	270	
ggc cac ctt cgt ggc ggg ctt cgt cgt ggg ctt cac cgc cgc ctg gca			864
Gly His Leu Arg Gly Gly Leu Arg Arg Gly Leu His Arg Arg Leu Ala			
275	280	285	
gct ggc gct cgt cac gct cgc cgt cgt gcc gct cat cgc cgt cat cgg			912
Ala Gly Ala Arg His Ala Arg Arg Arg Ala Ala His Arg Arg His Arg			

290	295	300	
ggg gct cag cgc cgc cgc gct cgc caa gct ctc ctc cag gag cca gga			960
Gly Ala Gln Arg Arg Arg Ala Arg Gln Ala Leu Leu Gln Glu Pro Gly			
305	310	315	320
cgc gct gtc ggg cgc cag cgg cat cgc gga gca ggc gct cgc gca gat			1008
Arg Ala Val Gly Arg Gln Arg His Arg Gly Ala Gly Ala Arg Ala Asp			
	325	330	335
acg gat cgt gca ggc ctt cgt cgg cga gga gcg cga aat gcg ggc gta			1056
Thr Asp Arg Ala Gly Leu Arg Arg Arg Gly Ala Arg Asn Ala Gly Val			
	340	345	350
ctc ggc ggc gtt ggc cgt cgc gca gaa gat cgg cta ccg cag cgg ctt			1104
Leu Gly Gly Val Gly Arg Arg Ala Glu Asp Arg Leu Pro Gln Arg Leu			
	355	360	365
cgc caa ggg gct cgg cct cgg cgg cac cta ctt cac cgt ctt ctg ctg			1152
Arg Gln Gly Ala Arg Pro Arg Arg His Leu Leu His Arg Leu Leu Leu			
	370	375	380
cta cgg cct cct gct ctg gta cgg cgg aca cct cgt ccg cgg aac cac			1200
Leu Arg Pro Pro Ala Leu Val Arg Arg Thr Pro Arg Pro Arg Asn His			
	385	390	400
acc aac gga ggg ctc gcc atc gcc acc atg ttc tcc gtc atg atc ggc			1248
Thr Asn Gly Gly Leu Ala Ile Ala Thr Met Phe Ser Val Met Ile Gly			
	405	410	415
ggg ctg gcc ctc ggg cag tcg gcg ccg agc atg gcc gcg ttc gcc aag			1296
Gly Leu Ala Leu Gly Gln Ser Ala Pro Ser Met Ala Ala Phe Ala Lys			
	420	425	430
gcg cgc gtg gcg gcc gcc aag atc ttc cgc atc atc gac cac agg ccg			1344
Ala Arg Val Ala Ala Ala Lys Ile Phe Arg Ile Ile Asp His Arg Pro			
	435	440	445
ggc atc tcc tcg cgg gac ggc gag gac ggc ggc ggc gtg gag ctg gag			1392
Gly Ile Ser Ser Arg Asp Gly Glu Asp Gly Gly Gly Val Glu Leu Glu			
	450	455	460
tcg gtg acg ggg cgg gtg gag atg agg ggc gtg gac ttc gcg tac ccg			1440
Ser Val Thr Gly Arg Val Glu Met Arg Gly Val Asp Phe Ala Tyr Pro			
	465	470	480
tcg cgg ccg gac gtc ccc atc ctg cgc ggc ttc tcg ctc agc gtg ccc			1488
Ser Arg Pro Asp Val Pro Ile Leu Arg Gly Phe Ser Leu Ser Val Pro			

485					490					495						
gcc	ggc	aag	acc	atc	gcg	ctg	gtg	ggc	agc	tcc	ggc	tcc	ggg	aag	agc	1536
Ala	Gly	Lys	Thr	Ile	Ala	Leu	Val	Gly	Ser	Ser	Gly	Ser	Gly	Lys	Ser	
			500					505					510			
acg	gtg	gtg	tcg	ctc	ctc	gag	agg	ttc	tac	gac	ccc	agc	gca	ggg	caa	1584
Thr	Val	Val	Ser	Leu	Leu	Glu	Arg	Phe	Tyr	Asp	Pro	Ser	Ala	Gly	Gln	
		515					520					525				
atc	ttg	ctg	gac	ggg	cat	gat	ctc	aag	tcg	ctg	aag	ctc	cgg	tgg	ctc	1632
Ile	Leu	Leu	Asp	Gly	His	Asp	Leu	Lys	Ser	Leu	Lys	Leu	Arg	Trp	Leu	
	530					535					540					
cgg	cag	cag	att	ggt	ctg	gtg	agc	cag	gag	ccg	acg	ctg	ttc	gcg	acg	1680
Arg	Gln	Gln	Ile	Gly	Leu	Val	Ser	Gln	Glu	Pro	Thr	Leu	Phe	Ala	Thr	
545					550					555					560	
agc	atc	aag	gag	aac	ctg	ctg	ctg	ggg	cgg	gac	agt	cag	agt	gcg	acg	1728
Ser	Ile	Lys	Glu	Asn	Leu	Leu	Leu	Gly	Arg	Asp	Ser	Gln	Ser	Ala	Thr	
				565					570					575		
cag	gcc	gag	atg	gag	gag	gcc	gcc	agg	gtg	gcc	aac	gcg	cac	tcc	ttc	1776
Gln	Ala	Glu	Met	Glu	Glu	Ala	Ala	Arg	Val	Ala	Asn	Ala	His	Ser	Phe	
			580					585					590			
atc	gtc	aag	ctc	ccc	gac	ggc	tac	gac	acg	cag	gtt	ggg	gag	cgc	ggc	1824
Ile	Val	Lys	Leu	Pro	Asp	Gly	Tyr	Asp	Thr	Gln	Val	Gly	Glu	Arg	Gly	
		595					600					605				
ctg	cag	ctc	tcc	ggt	ggg	cag	aag	cag	cgc	atc	gcc	atc	gcc	cgc	gcc	1872
Leu	Gln	Leu	Ser	Gly	Gly	Gln	Lys	Gln	Arg	Ile	Ala	Ile	Ala	Arg	Ala	
	610					615					620					
atg	ctc	aag	aac	ccc	gcc	atc	ctg	ctg	ctg	gac	gag	gcc	acc	agc	gcg	1920
Met	Leu	Lys	Asn	Pro	Ala	Ile	Leu	Leu	Leu	Asp	Glu	Ala	Thr	Ser	Ala	
625					630					635					640	
ctg	gac	tcc	gag	tct	gag	aag	ctc	gtg	cag	gag	gcg	ctg	gac	cgc	ttc	1968
Leu	Asp	Ser	Glu	Ser	Glu	Lys	Leu	Val	Gln	Glu	Ala	Leu	Asp	Arg	Phe	
				645					650					655		
atg	atc	ggg	cgc	acc	acc	ctg	gtg	atc	gcg	cac	agg	atg	tcc	acc	atc	2016
Met	Ile	Gly	Arg	Thr	Thr	Leu	Val	Ile	Ala	His	Arg	Met	Ser	Thr	Ile	
			660					665					670			
cgc	aag	gcc	gac	gtg	gtg	gcc	gtg	ctg	cag	ggc	ggc	ccc	gtc	tcc	gag	2064
Arg	Lys	Ala	Asp	Val	Val	Ala	Val	Leu	Gln	Gly	Gly	Pro	Val	Ser	Glu	

675	680	685	
atg ggc gcg cac gac gag ttg	atg gcc aag ggc gag aac ggc act tac		2112
Met Gly Ala His Asp Glu Leu	Met Ala Lys Gly Glu Asn Gly Thr Tyr		
690	695	700	
gcc aag ttc atc cgc atg cag gag cag gcg cac gag gcg gcg ttc gtc			2160
Ala Lys Phe Ile Arg Met Gln Glu Gln Ala His Glu Ala Ala Phe Val			
705	710	715	720
aac gcc cgc cgc agc agc gcc agg ccc tcc agc gcc cgc aac tcc gtc			2208
Asn Ala Arg Arg Ser Ser Ala Arg Pro Ser Ser Ala Arg Asn Ser Val			
	725	730	735
agc tcg ccc atc atg acg cgc aac tcc tcc tac ggc cgc tcc cca tac			2256
Ser Ser Pro Ile Met Thr Arg Asn Ser Ser Tyr Gly Arg Ser Pro Tyr			
	740	745	750
tcc cgc cgc ctc tcc gac ttc tcc acc tcc gac ttc acc ctc tcc atc			2304
Ser Arg Arg Leu Ser Asp Phe Ser Thr Ser Asp Phe Thr Leu Ser Ile			
	755	760	765
cac gac ccg cac cac cac cac cgg acg atg gcc gac aag cag ctc gcg			2352
His Asp Pro His His His His Arg Thr Met Ala Asp Lys Gln Leu Ala			
	770	775	780
ttc cgc gcc ggc gcc agc tcc ttc ctc cgc ctc gcc agg atg aac tcg			2400
Phe Arg Ala Gly Ala Ser Ser Phe Leu Arg Leu Ala Arg Met Asn Ser			
785	790	795	800
ccc gag tgg gcc tac gcg ctc gtc ggc tcc ctg ggc tcc atg gtc tgc			2448
Pro Glu Trp Ala Tyr Ala Leu Val Gly Ser Leu Gly Ser Met Val Cys			
	805	810	815
ggc tcc ttc agc gcc atc ttc gcc tac atc ctc agc gcc gtg ctc agc			2496
Gly Ser Phe Ser Ala Ile Phe Ala Tyr Ile Leu Ser Ala Val Leu Ser			
	820	825	830
gtc tac tac gcg ccg gac cct cgc tac atg aag cgc gag atc gcc aag			2544
Val Tyr Tyr Ala Pro Asp Pro Arg Tyr Met Lys Arg Glu Ile Ala Lys			
	835	840	845
tac tgc tac ctg ctc atc ggc atg tcc tcc gcg gcg ctg ctg ttc aac			2592
Tyr Cys Tyr Leu Leu Ile Gly Met Ser Ser Ala Ala Leu Leu Phe Asn			
850	855	860	
acg gtg cag cac gtg ttc tgg gac acg gtc ggc gag aac ctc acg aag			2640
Thr Val Gln His Val Phe Trp Asp Thr Val Gly Glu Asn Leu Thr Lys			

865	870	875	880	
cgt gtg cgc gag aag atg ttc gcc gcc gtg ctc cgc aac gag atc gcc				2688
Arg Val Arg Glu Lys Met Phe Ala Ala Val Leu Arg Asn Glu Ile Ala				
	885	890	895	
tgg ttc gac gcc gac gag aac gcc agc gcg cgc gtc gcc gcc agg ctc				2736
Trp Phe Asp Ala Asp Glu Asn Ala Ser Ala Arg Val Ala Ala Arg Leu				
	900	905	910	
gcg ctc gac gcc cag aac gtg cgc tcc gcc atc ggg gac cgt atc tcc				2784
Ala Leu Asp Ala Gln Asn Val Arg Ser Ala Ile Gly Asp Arg Ile Ser				
	915	920	925	
gtc atc gtc cag aac tcg gcg ctc atg ctc gtc gcc tgc acc gcg ggc				2832
Val Ile Val Gln Asn Ser Ala Leu Met Leu Val Ala Cys Thr Ala Gly				
	930	935	940	
ttc gtc ctc cag tgg cgc ctc gcg ctc gtg ctc ctc gcc gtc ttc ccg				2880
Phe Val Leu Gln Trp Arg Leu Ala Leu Val Leu Leu Ala Val Phe Pro				
	945	950	955	960
ctc gtc gtg gcc gcc acc gtg ctg cag aag atg ttc atg aag ggc ttc				2928
Leu Val Val Ala Ala Thr Val Leu Gln Lys Met Phe Met Lys Gly Phe				
	965	970	975	
tcg ggg gac ctg gag gcc gcg cac gcc agg gcc acg cag atc gcg ggc				2976
Ser Gly Asp Leu Glu Ala Ala His Ala Arg Ala Thr Gln Ile Ala Gly				
	980	985	990	
gag gcc gtg gcc aac ctg cgc acc gtg gcc gcg ttc aac gcg gag cgc				3024
Glu Ala Val Ala Asn Leu Arg Thr Val Ala Ala Phe Asn Ala Glu Arg				
	995	1000	1005	
aag atc acg ggg ctg ttc gag gcc aac ctg cgc ggc ccg ctc cgg cgc				3072
Lys Ile Thr Gly Leu Phe Glu Ala Asn Leu Arg Gly Pro Leu Arg Arg				
	1010	1015	1020	
tgc ttc tgg aag ggg cag atc gcc ggc agc ggc tac ggc gtg gcg cag				3120
Cys Phe Trp Lys Gly Gln Ile Ala Gly Ser Gly Tyr Gly Val Ala Gln				
	1025	1030	1035	1040
ttc ctg ctg tac gcg tcc tac gcg ctg ggg ctg tgg tac gcg gcg tgg				3168
Phe Leu Leu Tyr Ala Ser Tyr Ala Leu Gly Leu Trp Tyr Ala Ala Trp				
	1045	1050	1055	
ctg gtg aag cac ggc gtg tcc gac ttc tcg cgc acc atc cgc gtg ttc				3216
Leu Val Lys His Gly Val Ser Asp Phe Ser Arg Thr Ile Arg Val Phe				

1060	1065	1070	
atg gtg ctg atg gtg tcc gcc aac ggc gcc gcc gag acg ctg acg ctg			3264
Met Val Leu Met Val Ser Ala Asn Gly Ala Ala Glu Thr Leu Thr Leu			
1075	1080	1085	
gcg ccg gac ttt gtc aag ggc ggg cgc gcg atg cgg tcc gtg ttc gag			3312
Ala Pro Asp Phe Val Lys Gly Gly Arg Ala Met Arg Ser Val Phe Glu			
1090	1095	1100	
acc atc gac cgg aaa acg gag gtg gag ccc gac gac gtg gac gcg gcg			3360
Thr Ile Asp Arg Lys Thr Glu Val Glu Pro Asp Asp Val Asp Ala Ala			
1105	1110	1115	1120
ccg gtg ccg gag cgg ccc aag ggc gag gtg gag ctg aag cac gtg gac			3408
Pro Val Pro Glu Arg Pro Lys Gly Glu Val Glu Leu Lys His Val Asp			
1125	1130	1135	
ttc tcg tac ccg tcg cgg ccg gac atc cag gtg ttc cgc gac ctg agc			3456
Phe Ser Tyr Pro Ser Arg Pro Asp Ile Gln Val Phe Arg Asp Leu Ser			
1140	1145	1150	
ctc cgg gcg cgc gcc ggg aag acg ctg gcg ctg gtg ggt ccg agc ggg			3504
Leu Arg Ala Arg Ala Gly Lys Thr Leu Ala Leu Val Gly Pro Ser Gly			
1155	1160	1165	
tgc ggc aag agc tcg gtg ctg gcg ctg gtg cag cgg ttc tac gag ccc			3552
Cys Gly Lys Ser Ser Val Leu Ala Leu Val Gln Arg Phe Tyr Glu Pro			
1170	1175	1180	
acg tcc ggg cgc gtg ctc ctg gac ggc aag gac gtg cgc aag tac aac			3600
Thr Ser Gly Arg Val Leu Leu Asp Gly Lys Asp Val Arg Lys Tyr Asn			
1185	1190	1195	1200
ctg cgg gcg ctg cgg cgc gtg gtg gcg gtg gcg ccg cag gag ccg ttc			3648
Leu Arg Ala Leu Arg Arg Val Val Ala Val Ala Pro Gln Glu Pro Phe			
1205	1210	1215	
ctg ttc gcg gcg agc atc cac gac aac atc gcg tac ggg cgc gag ggc			3696
Leu Phe Ala Ala Ser Ile His Asp Asn Ile Ala Tyr Gly Arg Glu Gly			
1220	1225	1230	
gcg acg gag gcg gag gtg gtg gag gcg gcg acg cag gcg aac gcg cac			3744
Ala Thr Glu Ala Glu Val Val Glu Ala Ala Thr Gln Ala Asn Ala His			
1235	1240	1245	
cgg ttc atc gcg gcg ctg ccg gag ggc tac ggg acg cag gtg ggc gag			3792
Arg Phe Ile Ala Ala Leu Pro Glu Gly Tyr Gly Thr Gln Val Gly Glu			

1250	1255	1260	
cgc ggg gtg cag ctg tcg ggc ggg cag cgg cag cgg atc gcg atc gcg			3840
Arg Gly Val Gln Leu Ser Gly Gly Gln Arg Gln Arg Ile Ala Ile Ala			
1265	1270	1275	1280
cgc gcg ctg gtg aag cag gcg gcc atc gtg ctg ctg gac gag gcg acc			3888
Arg Ala Leu Val Lys Gln Ala Ala Ile Val Leu Leu Asp Glu Ala Thr			
1285	1290	1295	
agc gcg ctg gac gcc gag tcg gag cgg tgc gtg cag gag gcg ctg gag			3936
Ser Ala Leu Asp Ala Glu Ser Glu Arg Cys Val Gln Glu Ala Leu Glu			
1300	1305	1310	
cgc gcg ggg tcc ggg cgc acc acc atc gtg gtg gcg cac cgg ctg gcc			3984
Arg Ala Gly Ser Gly Arg Thr Thr Ile Val Val Ala His Arg Leu Ala			
1315	1320	1325	
acg gtg cgc ggc gcg cac acc atc gcg gtc atc gac gac ggc aag gtg			4032
Thr Val Arg Gly Ala His Thr Ile Ala Val Ile Asp Asp Gly Lys Val			
1330	1335	1340	
gcg gag cag ggg tcg cac tcg cac ctg ctc aag cac cat ccc gac ggg			4080
Ala Glu Gln Gly Ser His Ser His Leu Leu Lys His His Pro Asp Gly			
1345	1350	1355	1360
tgc tac gcg cgg atg ctg cag ctg cag cgg ctg acg ggc ggg tgc cgc			4128
Cys Tyr Ala Arg Met Leu Gln Leu Gln Arg Leu Thr Gly Gly Cys Arg			
1365	1370	1375	
gcc cgg gcc gcc gcc gtc gtc gtc caa cgg ggc cgc cgc gta gga tgg			4176
Ala Arg Ala Ala Ala Val Val Val Gln Arg Gly Arg Arg Val Gly Trp			
1380	1385	1390	
atg gat gga tca tgg atg agt ttg gtt cct tgataaa			4213
Met Asp Gly Ser Trp Met Ser Leu Val Pro			
1395	1400		

<210> 9

<211> 1402

<212> PRT

<213> Sorghum bicolor

<400> 9

Met Ser Thr Asn Asp Pro Asp Glu Ile Arg Ala Arg Val Val Val Leu

1

5

10

15

Gly	Ala	Pro	His	Ala	Asp	Asp	Asp	Ala	Gly	Asp	Glu	Trp	Ala	Arg	Pro	20	25	30
Glu	Leu	Glu	Ala	Phe	His	Leu	Pro	Ser	Pro	Ala	His	Gln	Pro	Pro	Gly	35	40	45
Phe	His	Leu	Ala	Ala	Gly	His	Gln	Pro	Glu	Ala	Ala	Ala	Glu	Gln	Pro	50	55	60
Thr	Thr	Leu	Pro	Ala	Ala	Arg	Arg	Thr	Ser	Asp	Thr	Ser	Thr	Ala	Ala	65	70	75
Gly	Ala	Ala	Pro	Pro	Ser	Pro	Ser	Pro	Pro	Pro	Pro	Pro	Ala	Pro	Leu	85	90	95
Glu	Met	Asp	Gln	Pro	Pro	Asn	Ala	Lys	Pro	Ala	Ser	Ser	Ser	Ala	Ala	100	105	110
Ala	Ala	Gly	Ala	Asn	Asp	Asn	Lys	Lys	Pro	Thr	Pro	Pro	Ala	Ala	Leu	115	120	125
Arg	Asp	Leu	Phe	Arg	Phe	Ala	Asp	Gly	Leu	Asp	Cys	Ala	Leu	Met	Leu	130	135	140
Val	Gly	Thr	Leu	Gly	Ala	Leu	Val	His	Gly	Cys	Ser	Leu	Pro	Val	Phe	145	150	155
Leu	Arg	Phe	Phe	Ala	Asp	Leu	Val	Asp	Ser	Phe	Gly	Ser	His	Ala	Asn	165	170	175
Asp	Pro	Asp	Thr	Met	Val	Arg	Leu	Val	Val	Lys	Tyr	Ala	Phe	Tyr	Phe	180	185	190
Leu	Val	Val	Gly	Ala	Ala	Ile	Trp	Ala	Ser	Ser	Trp	Ala	Glu	Ile	Ser	195	200	205
Cys	Trp	Met	Trp	Thr	Gly	Glu	Arg	Gln	Ser	Thr	Arg	Met	Arg	Ile	Arg	210	215	220
Tyr	Leu	Asp	Ala	Ala	Leu	Arg	Gln	Asp	Val	Ser	Phe	Phe	Asp	Thr	Asp	225	230	235
Val	Arg	Thr	Ser	Asp	Val	Ile	Tyr	Ala	Ile	Asn	Ala	Asp	Ala	Val	Val	245	250	255
Gly	Ala	Gly	Arg	His	Gln	Arg	Glu	Ala	Gly	Gln	Pro	His	Pro	Leu	His	260	265	270

Gly His Leu Arg Gly Gly Leu Arg Arg Gly Leu His Arg Arg Leu Ala
 275 280 285

Ala Gly Ala Arg His Ala Arg Arg Arg Ala Ala His Arg Arg His Arg
 290 295 300

Gly Ala Gln Arg Arg Arg Ala Arg Gln Ala Leu Leu Gln Glu Pro Gly
 305 310 315 320

Arg Ala Val Gly Arg Gln Arg His Arg Gly Ala Gly Ala Arg Ala Asp
 325 330 335

Thr Asp Arg Ala Gly Leu Arg Arg Arg Gly Ala Arg Asn Ala Gly Val
 340 345 350

Leu Gly Gly Val Gly Arg Arg Ala Glu Asp Arg Leu Pro Gln Arg Leu
 355 360 365

Arg Gln Gly Ala Arg Pro Arg Arg His Leu Leu His Arg Leu Leu Leu
 370 375 380

Leu Arg Pro Pro Ala Leu Val Arg Arg Thr Pro Arg Pro Arg Asn His
 385 390 395 400

Thr Asn Gly Gly Leu Ala Ile Ala Thr Met Phe Ser Val Met Ile Gly
 405 410 415

Gly Leu Ala Leu Gly Gln Ser Ala Pro Ser Met Ala Ala Phe Ala Lys
 420 425 430

Ala Arg Val Ala Ala Ala Lys Ile Phe Arg Ile Ile Asp His Arg Pro
 435 440 445

Gly Ile Ser Ser Arg Asp Gly Glu Asp Gly Gly Gly Val Glu Leu Glu
 450 455 460

Ser Val Thr Gly Arg Val Glu Met Arg Gly Val Asp Phe Ala Tyr Pro
 465 470 475 480

Ser Arg Pro Asp Val Pro Ile Leu Arg Gly Phe Ser Leu Ser Val Pro
 485 490 495

Ala Gly Lys Thr Ile Ala Leu Val Gly Ser Ser Gly Ser Gly Lys Ser
 500 505 510

Thr Val Val Ser Leu Leu Glu Arg Phe Tyr Asp Pro Ser Ala Gly Gln
 515 520 525

Ile	Leu	Leu	Asp	Gly	His	Asp	Leu	Lys	Ser	Leu	Lys	Leu	Arg	Trp	Leu		
530						535					540						
Arg	Gln	Gln	Ile	Gly	Leu	Val	Ser	Gln	Glu	Pro	Thr	Leu	Phe	Ala	Thr		
545					550					555					560		
Ser	Ile	Lys	Glu	Asn	Leu	Leu	Leu	Gly	Arg	Asp	Ser	Gln	Ser	Ala	Thr		
				565					570					575			
Gln	Ala	Glu	Met	Glu	Glu	Ala	Ala	Arg	Val	Ala	Asn	Ala	His	Ser	Phe		
			580					585					590				
Ile	Val	Lys	Leu	Pro	Asp	Gly	Tyr	Asp	Thr	Gln	Val	Gly	Glu	Arg	Gly		
		595					600					605					
Leu	Gln	Leu	Ser	Gly	Gly	Gln	Lys	Gln	Arg	Ile	Ala	Ile	Ala	Arg	Ala		
	610					615					620						
Met	Leu	Lys	Asn	Pro	Ala	Ile	Leu	Leu	Leu	Asp	Glu	Ala	Thr	Ser	Ala		
625					630					635					640		
Leu	Asp	Ser	Glu	Ser	Glu	Lys	Leu	Val	Gln	Glu	Ala	Leu	Asp	Arg	Phe		
				645					650					655			
Met	Ile	Gly	Arg	Thr	Thr	Leu	Val	Ile	Ala	His	Arg	Met	Ser	Thr	Ile		
			660					665					670				
Arg	Lys	Ala	Asp	Val	Val	Ala	Val	Leu	Gln	Gly	Gly	Pro	Val	Ser	Glu		
		675					680					685					
Met	Gly	Ala	His	Asp	Glu	Leu	Met	Ala	Lys	Gly	Glu	Asn	Gly	Thr	Tyr		
	690					695					700						
Ala	Lys	Phe	Ile	Arg	Met	Gln	Glu	Gln	Ala	His	Glu	Ala	Ala	Phe	Val		
705					710					715					720		
Asn	Ala	Arg	Arg	Ser	Ser	Ala	Arg	Pro	Ser	Ser	Ala	Arg	Asn	Ser	Val		
				725				730						735			
Ser	Ser	Pro	Ile	Met	Thr	Arg	Asn	Ser	Ser	Tyr	Gly	Arg	Ser	Pro	Tyr		
			740					745					750				
Ser	Arg	Arg	Leu	Ser	Asp	Phe	Ser	Thr	Ser	Asp	Phe	Thr	Leu	Ser	Ile		
		755					760					765					
His	Asp	Pro	His	His	His	His	Arg	Thr	Met	Ala	Asp	Lys	Gln	Leu	Ala		
	770					775					780						

Phe Arg Ala Gly Ala Ser Ser Phe Leu Arg Leu Ala Arg Met Asn Ser
785 790 795 800

Pro Glu Trp Ala Tyr Ala Leu Val Gly Ser Leu Gly Ser Met Val Cys
805 810 815

Gly Ser Phe Ser Ala Ile Phe Ala Tyr Ile Leu Ser Ala Val Leu Ser
820 825 830

Val Tyr Tyr Ala Pro Asp Pro Arg Tyr Met Lys Arg Glu Ile Ala Lys
835 840 845

Tyr Cys Tyr Leu Leu Ile Gly Met Ser Ser Ala Ala Leu Leu Phe Asn
850 855 860

Thr Val Gln His Val Phe Trp Asp Thr Val Gly Glu Asn Leu Thr Lys
865 870 875 880

Arg Val Arg Glu Lys Met Phe Ala Ala Val Leu Arg Asn Glu Ile Ala
885 890 895

Trp Phe Asp Ala Asp Glu Asn Ala Ser Ala Arg Val Ala Ala Arg Leu
900 905 910

Ala Leu Asp Ala Gln Asn Val Arg Ser Ala Ile Gly Asp Arg Ile Ser
915 920 925

Val Ile Val Gln Asn Ser Ala Leu Met Leu Val Ala Cys Thr Ala Gly
930 935 940

Phe Val Leu Gln Trp Arg Leu Ala Leu Val Leu Leu Ala Val Phe Pro
945 950 955 960

Leu Val Val Ala Ala Thr Val Leu Gln Lys Met Phe Met Lys Gly Phe
965 970 975

Ser Gly Asp Leu Glu Ala Ala His Ala Arg Ala Thr Gln Ile Ala Gly
980 985 990

Glu Ala Val Ala Asn Leu Arg Thr Val Ala Ala Phe Asn Ala Glu Arg
995 1000 1005

Lys Ile Thr Gly Leu Phe Glu Ala Asn Leu Arg Gly Pro Leu Arg Arg
1010 1015 1020

Cys Phe Trp Lys Gly Gln Ile Ala Gly Ser Gly Tyr Gly Val Ala Gln
1025 1030 1035 1040

Phe Leu Leu Tyr Ala Ser Tyr Ala Leu Gly Leu Trp Tyr Ala Ala Trp
1045 1050 1055

Leu Val Lys His Gly Val Ser Asp Phe Ser Arg Thr Ile Arg Val Phe
1060 1065 1070

Met Val Leu Met Val Ser Ala Asn Gly Ala Ala Glu Thr Leu Thr Leu
1075 1080 1085

Ala Pro Asp Phe Val Lys Gly Gly Arg Ala Met Arg Ser Val Phe Glu
1090 1095 1100

Thr Ile Asp Arg Lys Thr Glu Val Glu Pro Asp Asp Val Asp Ala Ala
1105 1110 1115 1120

Pro Val Pro Glu Arg Pro Lys Gly Glu Val Glu Leu Lys His Val Asp
1125 1130 1135

Phe Ser Tyr Pro Ser Arg Pro Asp Ile Gln Val Phe Arg Asp Leu Ser
1140 1145 1150

Leu Arg Ala Arg Ala Gly Lys Thr Leu Ala Leu Val Gly Pro Ser Gly
1155 1160 1165

Cys Gly Lys Ser Ser Val Leu Ala Leu Val Gln Arg Phe Tyr Glu Pro
1170 1175 1180

Thr Ser Gly Arg Val Leu Leu Asp Gly Lys Asp Val Arg Lys Tyr Asn
1185 1190 1195 1200

Leu Arg Ala Leu Arg Arg Val Val Ala Val Ala Pro Gln Glu Pro Phe
1205 1210 1215

Leu Phe Ala Ala Ser Ile His Asp Asn Ile Ala Tyr Gly Arg Glu Gly
1220 1225 1230

Ala Thr Glu Ala Glu Val Val Glu Ala Ala Thr Gln Ala Asn Ala His
1235 1240 1245

Arg Phe Ile Ala Ala Leu Pro Glu Gly Tyr Gly Thr Gln Val Gly Glu
1250 1255 1260

Arg Gly Val Gln Leu Ser Gly Gly Gln Arg Gln Arg Ile Ala Ile Ala
1265 1270 1275 1280

Arg Ala Leu Val Lys Gln Ala Ala Ile Val Leu Leu Asp Glu Ala Thr
1285 1290 1295

Ser Ala Leu Asp Ala Glu Ser Glu Arg Cys Val Gln Glu Ala Leu Glu
1300 1305 1310

Arg Ala Gly Ser Gly Arg Thr Thr Ile Val Val Ala His Arg Leu Ala
1315 1320 1325

Thr Val Arg Gly Ala His Thr Ile Ala Val Ile Asp Asp Gly Lys Val
1330 1335 1340

Ala Glu Gln Gly Ser His Ser His Leu Leu Lys His His Pro Asp Gly
1345 1350 1355 1360

Cys Tyr Ala Arg Met Leu Gln Leu Gln Arg Leu Thr Gly Gly Cys Arg
1365 1370 1375

Ala Arg Ala Ala Ala Val Val Val Gln Arg Gly Arg Arg Val Gly Trp
1380 1385 1390

Met Asp Gly Ser Trp Met Ser Leu Val Pro
1395 1400

Sequence